

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90092 A1

(51) International Patent Classification⁷: C07D 277/52, 277/60, A61K 31/426, A61P 3/00, 5/48, 27/06, 25/24, 25/28, 29/00, 31/12, 31/06

(74) Agents: BERG, Sven, Anders et al.; Albihns Stockholm AB, Box 5581, S-114 85 Stockholm (SE).

(21) International Application Number: PCT/SE01/01158

(22) International Filing Date: 22 May 2001 (22.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0001899-4 22 May 2000 (22.05.2000) SE

(71) Applicant (for all designated States except US): BIOVIT-RUM AB [SE/SE]; S-112 76 Stockholm (SE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/90092 A1

(72) Inventors; and
(75) Inventors/Applicants (for US only): KURZ, Guido [DE/SE]; Möregatan 10, 6tr, S-118 27 Stockholm (SE). NILSSON, Marianne [SE/SE]; PI 2654 Ubbby, S-762 94 Rimbo (SE).

(54) Title: INHIBITORS OF 11-BETA-HYDROXY STEROID DEHYDROGENASE TYPE 1

(57) Abstract: The present invention relates to compounds with the formula (I) and also to pharmaceutical compositions comprising the compounds, to processes for their preparation, as well as to the use of the compounds in medicine and for the preparation of a medicament which acts on the human 11- β -hydroxysteroid dehydrogenase type 1 enzyme.

INHIBITORS OF 11-BETA-HYDROXY STEROID DEHYDROGENASE TYPE 1

5 TECHNICAL FIELD

The present invention relates to novel compounds, to pharmaceutical compositions comprising the compounds, to processes for their preparation, as well as to the use of the compounds in medicine and for the preparation of a medicament which acts on the
10 human 11- β -hydroxysteroid dehydrogenase type 1 enzyme (11 β HSD1).

BACKGROUND ART

1. Glucocorticoids, diabetes and hepatic glucose production

15

It has been known for more than half a century that glucocorticoids have a central role in diabetes, e.g. the removal of the pituitary or the adrenal gland from a diabetic animal alleviates the most severe symptoms of diabetes and lowers the concentration of glucose in the blood (Long, C.D. and F.D.W. Leukins (1936) J. Exp. Med. 63: 465-
20 490; Houssay, B.A. (1942) Endocrinology 30: 884-892). It is also well established that glucocorticoids enable the effect of glucagon on the liver.

The role of 11 β HSD1 as an important regulator of local glucocorticoid effect and thus of hepatic glucose production is well substantiated (see e.g. Jamieson et al. (2000) J.
25 Endocrinol. 165: p. 685-692). The hepatic insulin sensitivity was improved in healthy human volunteers treated with the non-specific 11 β HSD1 inhibitor carbenoxolone (Walker, B.R. et al. (1995) J. Clin. Endocrinol. Metab. 80: 3155-3159). Furthermore, the expected mechanism has been established by different experiments with mice and rats. These studies showed that the mRNA levels and activities of two key enzymes in
30 hepatic glucose production were reduced, namely: the rate-limiting enzyme in gluconeogenesis, phosphoenolpyruvate carboxykinase (PEPCK), and glucose-6-

phosphatase (G6Pase) catalyzing the last common step of gluconeogenesis and glycogenolysis. Finally, the blood glucose level and hepatic glucose production is reduced in mice having the 11β HSD1 gene knocked-out. Data from this model also confirm that inhibition of 11β HSD1 will not cause hypoglycemia, as predicted since the basal levels of PEPCCK and G6Pase are regulated independently of glucocorticoids (Kotelevtsev, Y. et al., (1997) Proc. Natl. Acad. Sci. USA 94: 14924-14929).

2. Possible reduction of obesity and obesity related cardiovascular risk factors

Obesity is an important factor in syndrome X as well as in the majority (> 80%) of type 2 diabetic, and omental fat appears to be of central importance. Abdominal obesity is closely associated with glucose intolerance, hyperinsulinemia, hypertriglyceridemia, and other factors of the so-called syndrome X (e.g. raised blood pressure, decreased levels of HDL and increased levels of VLDL) (Montague & O'Rahilly, Diabetes 49: 883-888, 2000). Inhibition of the enzyme in pre-adipocytes (stromal cells) has been shown to decrease the rate of differentiation into adipocytes. This is predicted to result in diminished expansion (possibly reduction) of the omental fat depot, i.e. reduced central obesity (Bujalska, I.J., S. Kumar, and P.M. Stewart (1997) Lancet 349: 1210-1213).

Inhibition of 11β HSD1 in mature adipocytes is expected to attenuate secretion of the plasminogen activator inhibitor 1 (PAI-1) – an independent cardiovascular risk factor (Halleux, C.M. et al. (1999) J. Clin. Endocrinol. Metab. 84: 4097-4105). Furthermore, there is a clear correlation between glucocorticoid “activity” and cardiovascular risk factors suggesting that a reduction of the glucocorticoid effects would be beneficial (Walker, B.R. et al. (1998) Hypertension 31: 891-895; Fraser, R. et al. (1999) Hypertension 33: 1364-1368).

Adrenalectomy attenuates the effect of fasting to increase both food intake and hypothalamic neuropeptide Y expression. This supports the role of glucocorticoids in promoting food intake and suggests that inhibition of 11β HSD1 in the brain might

increase satiety and therefore reduce food intake (Woods, S.C. et al. (1998) Science, 280: 1378-1383).

3. Possible beneficial effect on the pancreas

5

Inhibition of 11 β HSD1 in isolated murine pancreatic β -cells improves the glucose-stimulated insulin secretion (Davani, B. et al. (2000) J. Biol. Chem. 2000 Nov 10; 275(45): 34841-4). Glucocorticoids were previously known to reduce pancreatic insulin release *in vivo* (Billaudel, B. and B.C.J. Sutter (1979) Horm. Metab. Res. 11: 555-560). Thus, inhibition of 11 β HSD1 is predicted to yield other beneficial effects for diabetes treatment, besides effects on liver and fat.

4. Possible beneficial effects on cognition and dementia

15 Stress and glucocorticoids influence cognitive function (de Quervain, D.J.-F., B. Roozendaal, and J.L. McGaugh (1998) Nature 394: 787-790). The enzyme 11 β HSD1 controls the level of glucocorticoid action in the brain and thus contributes to neurotoxicity (Rajan, V., C.R.W. Edwards, and J.R. Seckl, J. (1996) Neuroscience 16: 65-70; Seckl, J.R., Front. (2000) Neuroendocrinol. 18: 49-99). Unpublished results
20 indicate significant memory improvement in rats treated with a non-specific 11 β HSD1 inhibitor (J. Seckl, personal communication). Based the above and on the known effects of glucocorticoids in the brain, it may also be suggested that inhibiting 11 β HSD1 in the brain may result in reduced anxiety (Tronche, F. et al. (1999) Nature Genetics 23: 99-103). Thus, taken together, the hypothesis is that inhibition of
25 11 β HSD1 in the human brain would prevent reactivation of cortisone into cortisol and protect against deleterious glucocorticoid-mediated effects on neuronal survival and other aspects of neuronal function, including cognitive impairment, depression, and increased appetite (previous section).

5. Possible use of immuno-modulation using 11 β HSD1 inhibitors

The general perception is that glucocorticoids suppress the immune system. But in fact there is a dynamic interaction between the immune system and the HPA
5 (hypothalamo-pituitary-adrenal) axis (Rook, G.A.W. (1999) Baillière's Clin. Endocrinol. Metab. 13: 576-581). The balance between the cell-mediated response and humoral responses is modulated by glucocorticoids. A high glucocorticoid activity, such as at a state of stress, is associated with a humoral response. Thus, inhibition of the enzyme 11 β HSD1 has been suggested as a means of shifting the response towards
10 a cell-based reaction.

In certain disease states, including tuberculosis, lepra and psoriasis the immune reaction is normally biased towards a humoral response when in fact the appropriate response would be cell based. Temporal inhibition of 11 β HSD1, local or systemic,
15 might be used to push the immune system into the appropriate response (Mason, D. (1991) Immunology Today 12: 57-60; Rook et al., *supra*).

An analogous use of 11 β HSD1 inhibition, in this case temporal, would be to booster the immune response in association with immunization to ensure that a cell based
20 response would be obtained, when desired.

6. Reduction of intraocular pressure

Recent data suggest that the levels of the glucocorticoid target receptors and the
25 11 β HSD enzymes determines the susceptibility to glaucoma (Stokes, J. et al. (2000) Invest. Ophthalmol. 41: 1629-1638). Further, inhibition of 11 β HSD1 was recently presented as a novel approach to lower the intraocular pressure (Walker E. A. et al, poster P3-698 at the Endocrine society meeting June 12-15, 1999, San Diego). Ingestion of carbenoxolone, a non-specific inhibitor of 11 β HSD1, was shown to
30 reduce the intraocular pressure by 20% in normal subjects. In the eye, expression of 11 β HSD1 is confined to basal cells of the corneal epithelium and the non-pigmented

epithelium of the cornea (the site of aqueous production), to ciliary muscle and to the sphincter and dilator muscles of the iris. In contrast, the distant isoenzyme 11 β HSD2 is highly expressed in the non-pigmented ciliary epithelium and corneal endothelium. None of the enzymes is found at the trabecular meshwork, the site of drainage. Thus, 11 β HSD1 is suggested to have a role in aqueous production, rather than drainage, but it is presently unknown if this is by interfering with activation of the glucocorticoid or the mineralocorticoid receptor, or both.

7. Reduced osteoporosis

10

Glucocorticoids have an essential role in skeletal development and function but are detrimental in excess. Glucocorticoid-induced bone loss is derived, at least in part, via inhibition of bone formation, which includes suppression of osteoblast proliferation and collagen synthesis (Kim, C.H., S.L. Cheng, and G.S. Kim (1999) J. Endocrinol. 162: 371-379). The negative effect on bone nodule formation could be blocked by the non-specific inhibitor carbenoxolone suggesting an important role of 11 β HSD1 in the glucocorticoid effect (Bellows, C.G., A. Ciaccia, and J.N.M. Heersche, (1998) Bone 23: 119-125). Other data suggest a role of 11 β HSD1 in providing sufficiently high levels of active glucocorticoid in osteoclasts, and thus in augmenting bone resorption (Cooper, M.S. et al. (2000) Bone 27: 375-381). Taken together, these different data suggest that inhibition of 11 β HSD1 may have beneficial effects against osteoporosis by more than one mechanism working in parallel.

WO 99/65884 discloses carbon substituted aminothiazole inhibitors of cyclin dependent kinases. These compounds may e.g. be used against cancer, inflammation and arthritis. US 5,856,347 discloses an antibacterial preparation or bactericide comprising 2-aminothiazole derivative and/or salt thereof. Further, US 5,403,857 discloses benzenesulfonamide derivatives having 5-lipoxygenase inhibitory activity. Additionally, tetrahydrothiazolo[5,4-c]pyridines are disclosed in: Analgesic tetrahydrothiazolo[5,4-c]pyridines. Fr. Addn. (1969), 18 pp, Addn. to Fr. 1498465. CODEN: FAXXA3; FR 94123 19690704 CAN 72:100685 AN 1970:100685 CAPLUS

and 4,5,6,7-Tetrahydrothiazolo[5,4-c]pyridines. Neth. Appl. (1967), 39 pp. CODEN: NAXXAN NL 6610324 19670124 CAN 68:49593, AN 1968: 49593 CAPLUS.

FR 2384498 discloses thiazolo-benzenesulfonamides which show antibacterial,
5 antifungal and hypoglycaemic properties. WO99/28306 and EP 0 819 681 A2 relate to thiazolobenzenesulfonamides which can be used for treating neurodegenerative pathologies, such as Alzheimer's disease. JP 7149745 A2 and JP 7149746 A2 both describe 2-aminothiazole derivatives as esterase inhibitors. Nothing is disclosed about inhibiting 11 β HSD1. JP 7309757 A2 relates to treating Alzheimer's disease using N-
10 (5-nitro-2-thiazolyl)benzenesulfonamides. JP 3173876 A2 presents preparation of diphenylthiazoles. These compounds are used as anti-inflammatories, analgesics, anti-allergy agents, uric acid accelerators and blood platelet aggregation inhibitors. EP 0 790 057 A1 discloses an antibacterial or bactericide comprising a 2-aminothiazole derivative. US 2 362 087 describes the preparation of thiazolobenzenesulfonamides,
15 such as 2-bromobenzenesulfonamido-4-methylthiazole. Nothing is disclosed about inhibiting 11 β HSD1 and no therapeutic use of such substances is disclosed.

However, none of the above disclosures discloses the compounds according to the present invention, or their use for the treatment of diabetes, obesity, glaucoma,
20 osteoporosis, cognitive disorders, immune disorders, and depression.


Consequently, there is a need of new compounds that are useful in the treatment of diabetes, obesity, glaucoma, osteoporosis, cognitive disorders, immune disorders, and depression.
25

DISCLOSURE OF THE INVENTION

The compounds according to the present invention solve the above problems and embraces a novel class of compounds which has been developed and which inhibit the
30 human 11- β -hydroxysteroid dehydrogenase type 1 enzyme (11- β -HSD₁), and may

therefore be of use in the treating disorders such as diabetes, obesity, glaucoma, osteoporosis, cognitive disorders; immune disorders, and depression.

One object of the present invention is A compound of the formula (I):



5

wherein

T is selected from

- thienyl substituted with one or more of bromo, chloro;
 10 phenyl substituted as follows:

- a) either T is phenyl, wherein the phenyl is substituted with one or more of propyl and phenyl;
- b) T is phenyl substituted with chloro in position 3 and methyl in position 2;
- 15 c) T is phenyl substituted with chloro in position 2 and 4, and methyl in position 6;
- d) T is phenyl substituted with bromo in position 4 and fluoro in position 2 and 5;
- e) T is phenyl substituted with chloro in position 2, 3, and 4;
- f) T is phenyl substituted with chloro in position 2, 4, and 5;
- g) T is phenyl substituted with bromo in position 4 and methyl in position 2;
- 20 h) T is phenyl substituted with chloro in position 2 and 6;
- i) T is phenyl substituted with chloro in position 2, 4, and 6; or
- j) T is phenyl substituted with bromo in position 4 and chloro in position 5.

- A is selected from an aryl ring or heteroaryl ring, which can further be optionally
 25 substituted in one or more positions independently of each other by hydrogen, C₁₋₆-alkyl, halogenated C₁₋₆-alkyl, halogen, C₁₋₆-alkoxy, nitro, C₁₋₆-alkoxycarbonyl, C₁₋₆-alkylsulfonyl, acetylamino or aryloxy, wherein the aryloxy can further be optionally

substituted in one or more positions independently of each other by hydrogen and halogen;

- 5 B is selected from hydrogen and C₁₋₆-alkoxycarbonyl or is linked to A to give a 6-membered aromatic or non-aromatic ring;

as well as pharmaceutically acceptable salts, hydrates and solvates thereof.

It is preferred that:

- 10 A is selected from 1-benzothien-3-yl, 3-(2,5-dimethylfuryl), pyridinyl; thienyl optionally substituted with one or more of chloro, methylsulfonyl; phenyl optionally substituted with one or more of ethoxycarbonyl, nitro, fluoro, methyl, methoxy, acetylamino, chloro, 4-chlorophenoxy, trifluoromethyl;
- 15 B is selected from hydrogen, carbethoxy or is linked to A to give a 6-membered aromatic or non-aromatic ring.

Specific examples of compounds according to the present invention are:

- 20 Ethyl 2-[[[(3-chloro-2-methylphenyl)sulfonyl]amino]-4-phenyl-1,3-thiazole-5-carboxylate,
- N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- N-(4-phenyl-1,3-thiazol-2-yl)-4-propylbenzenesulfonamide,
- N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- 25 3-Chloro-2-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 3-Chloro-2-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
- 3-Chloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 30 2,4,6-Trichloro-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,

- 2,4,6-Trichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-(4-phenyl-1,3-thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Fluoro-3-methylphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
5 N-[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
2,4-Dichloro-6-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4-Dichloro-6-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4-Dichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-6-
methylbenzenesulfonamide,
10 2,4-Dichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
N-[4-(2-{{[(4-propylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl})phenyl]acetamide,
4-Propyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-
15 propylbenzenesulfonamide,
N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-{{[(3-chloro-2-methylphenyl)sulfonyl]amino}-1,3-thiazol-4-
yl})phenyl]acetamide,
20 3-Chloro-2-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-2-
methylbenzenesulfonamide,
3-Chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-
methylbenzenesulfonamide,
25 3-Chloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
3-Chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
N-[4-(2-{{[(2,4,6-trichlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl})phenyl]acetamide,
2,4,6-Trichloro-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
30 2,4,6-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-
yl)benzenesulfonamide,

- 2,4,6-Trichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-(4-{2-[(1,1'-biphenyl)-4-ylsulfonyl]amino}-1,3-thiazol-4-yl)phenylacetamide,
N-[4-(3-pyridinyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
5 N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
10 N-[4-(2-{[(2,4-dichloro-6-methylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
2,4-Dichloro-6-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4-Dichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
15 2,4-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-6-methylbenzenesulfonamide,
2,4-Dichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
2,4-Dichloro-N-[4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
20 N-[4-(1-benzothien-3-yl)-1,3-thiazol-2-yl]-2,4-dichloro-6-methylbenzenesulfonamide,
N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
3-Chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
2,4,6-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
25 2,4-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
2,4-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
Ethyl 2-[(1,1'-biphenyl)-4-ylsulfonyl]amino-4-phenyl-1,3-thiazole-5-carboxylate,
3-Chloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
30 N-[4-(2-{[(4-Bromo-2,5-difluorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,

- 2,3,4-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,5-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,3,4-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
4-Bromo-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2,5-difluorobenzenesulfonamide,
5 4,5-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
4,5-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
N-[4-(2-((2,4,5-Trichlorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl]phenyl]acetamide,
2,3,4-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
10 4-Bromo-5-chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
3-Bromo-5-chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
15 3-Bromo-5-chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
N-[4-(2-((2,6-Dichlorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl]phenyl]acetamide,
2,6-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-(7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
20 2,3,4-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
N-[4-(2-Chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
25 4-Bromo-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
4-Bromo-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
4,5-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
30

- 4-Bromo-5-chloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 4-Bromo-5-chloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 5 2,4-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 4-Bromo-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-
- 10 methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- N-[4-(2-{[(4-Bromo-5-chloro-2-thienyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
- 15 3-Chloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- 20 2,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-6-methylbenzenesulfonamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-4-propylbenzenesulfonamide,
- 3-Chloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-2-methylbenzenesulfonamide,
- 25 N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-4-propylbenzenesulfonamide,
- 2,4-Dichloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-6-methylbenzenesulfonamide,
- N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl][1,1'-
- 30 biphenyl]-4-sulfonamide,

N-(2-{{[(3-Chloro-2-methylphenyl)sulfonyl]amino}naphtho[1,2-d][1,3]thiazol-6-yl)acetamide,

N-(2-{{[(4-Propylphenyl)sulfonyl]amino}-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl)acetamide,

- 5 N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide,
N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide.

Another object of the present invention is a compound as described above for medical use.


10

The compounds described above can be prepared by methods comprising at least one of the following steps:

- a) sulfonamide coupling by reacting a 2-aminothiazole with a sulfonylchloride in the presence of a base,
- 15 b) sulfonamide coupling by reacting a 2-aminothiazole derivative with a sulfonylchloride in the presence of a base,
- c) formation of a thiazole ring by reacting an optionally substituted thiourea with an α -haloketone,
- d) formation of a thiazole ring by reacting a thiourea with a ketone.

20

- Another object of the present invention is a method for the treatment or prevention of diabetes, syndrome X, obesity, glaucoma, hyperlipidemia, hyperglycemia, hyperinsulinemia, osteoporosis, tuberculosis, depression, virus diseases and inflammatory disorders, said method comprising administering to a mammal,
- 25 including man, in need of such treatment an effective amount of a compound of the formula (I)



wherein

T is an aryl ring or heteroaryl ring, optionally independently substituted by $[R]_n$,
 5 wherein n is an integer 0-5, and R is hydrogen, halogen, C_{1-6} -alkyl, and aryl;

A is selected from an aryl ring or heteroaryl ring, which can further be optionally substituted in one or more positions independently of each other by hydrogen, C_{1-6} -alkyl, halogenated C_{1-6} -alkyl, halogen, C_{1-6} -alkoxy, nitro, C_{1-6} -alkoxycarbonyl, C_{1-6} -alkylsulfonyl, acetylamino or aryloxy, wherein the aryloxy can further be optionally substituted in one or more positions independently of each other by hydrogen and halogen;
 10

B is selected from hydrogen and C_{1-6} -alkoxycarbonyl or is linked to A to give a 6-membered aromatic or non-aromatic ring;
 15

as well as pharmaceutically acceptable salts, hydrates and solvates thereof.

These compounds may also be used in the in the manufacture of a medicament for the
 20 treatment or prevention of diabetes, syndrome X, obesity, glaucoma, hyperlipidemia, hyperglycemia, hyperinsulinemia, osteoporosis, tuberculosis, depression, virus diseases and inflammatory disorders.

It is preferred that:

25 T is selected from
 thienyl substituted with one or more of bromo, chloro;

phenyl optionally substituted with one or more of chloro, methyl, propyl, phenyl, bromo, fluoro;

A is selected from 1-benzothien-3-yl, 3-(2,5-dimethylfuryl), pyridinyl;
5 thienyl optionally substituted with one or more of chloro, methylsulfonyl;
phenyl optionally substituted with one or more of ethoxycarbonyl, nitro, fluoro, methyl, methoxy, acetylamino, chloro, 4-chlorophenoxy, trifluoromethyl;

10 B is selected from hydrogen, carbethoxy or is linked to A to give a 6-membered aromatic or non-aromatic ring.

Specific examples of compounds according to the present invention are:

Ethyl 2-(2-{[(4-chlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)benzoate,
2,5-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
15 4-Chloro-N-[4-(4,5-dichloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
Ethyl 2-{[(4-chlorophenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-carboxylate,
Ethyl 2-{[(3-chloro-2-methylphenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-carboxylate,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
20 N-(4-phenyl-1,3-thiazol-2-yl)-4-propylbenzenesulfonamide,
N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
3-Chloro-2-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-2-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
25 3-Chloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
2,4,6-Trichloro-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4,6-Trichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
30 2,4,6-Trichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,

- N-(4-phenyl-1,3-thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Fluoro-3-methylphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
2,4-Dichloro-6-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
5 2,4-Dichloro-6-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4-Dichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
2,4-Dichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
N-[4-(2-{{[(4-propylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl})phenyl]acetamide,
10 4-Propyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide,
N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
15 N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-{{[(3-chloro-2-methylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl})phenyl]acetamide,
3-Chloro-2-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
20 3-Chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-methylbenzenesulfonamide,
3-Chloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
3-Chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
25 N-[4-(2-{{[(2,4,6-trichlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl})phenyl]acetamide,
2,4,6-Trichloro-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
30 2,4,6-Trichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,

- N-(4-{2-[(1,1'-biphenyl)-4-ylsulfonyl]amino}-1,3-thiazol-4-yl)phenylacetamide,
 N-[4-(3-pyridinyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
 N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
 N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
 5 N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
 N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
 N-[4-(2-{[(2,4-dichloro-6-methylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
 10 2,4-Dichloro-6-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
 2,4-Dichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
 2,4-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-6-methylbenzenesulfonamide,
 15 2,4-Dichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
 2,4-Dichloro-N-[4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
 N-[4-(1-benzothien-3-yl)-1,3-thiazol-2-yl]-2,4-dichloro-6-methylbenzenesulfonamide,
 20 N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
 3-Chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
 2,4,6-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
 2,4-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
 25 2,4-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
 Ethyl 2-[(1,1'-biphenyl)-4-ylsulfonyl]amino-4-phenyl-1,3-thiazole-5-carboxylate,
 3-Chloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
 N-[4-(2-{[(4-Bromo-2,5-difluorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
 30 2,3,4-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
 2,4,5-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,

- 2,3,4-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
4-Bromo-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2,5-difluorobenzenesulfonamide,
4,5-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-
thiophenesulfonamide,
- 5 4,5-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
N-[4-(2-((2,4,5-Trichlorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl)phenyl]acetamide,
2,3,4-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-
yl)benzenesulfonamide,
4-Bromo-5-chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-
- 10 thiophenesulfonamide,
3-Bromo-5-chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-
thiophenesulfonamide,
3-Bromo-5-chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
N-[4-(2-((2,6-Dichlorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl)phenyl]acetamide,
- 15 2,6-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-(7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-
yl)benzenesulfonamide,
2,3,4-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-
yl}benzenesulfonamide,
- 20 2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-
yl}benzenesulfonamide,
N-[4-(2-Chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
4-Bromo-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2,5-
difluorobenzenesulfonamide,
- 25 4-Bromo-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2,5-
difluorobenzenesulfonamide,
4,5-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-
thiophenesulfonamide,
4-Bromo-5-chloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2-
- 30 thiophenesulfonamide,

- 4-Bromo-5-chloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 5 2,4,6-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 4-Bromo-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 10 N-[4-(2-[(4-Bromo-5-chloro-2-thienyl)sulfonyl]amino)-1,3-thiazol-4-yl]phenyl]acetamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 3,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 15 3-Chloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- 20 2,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-6-methylbenzenesulfonamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-4-propylbenzenesulfonamide,
- 3-Chloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-2-methylbenzenesulfonamide,
- 25 N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-4-propylbenzenesulfonamide,
- 2,4-Dichloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-6-methylbenzenesulfonamide,
- N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- 30 N-{2-[(Phenylsulfonyl)amino]-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide,

- N-(2-{{[(3-Chloro-2-methylphenyl)sulfonyl]amino}naphtho[1,2-d][1,3]thiazol-6-yl)acetamide,
- N-(2-{{[(4-Propylphenyl)sulfonyl]amino}-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl)acetamide,
- 5 N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide,
N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide.

Another object of the present invention is a pharmaceutical composition comprising at
10 least one compound of the formula (I) as defined above, and a pharmaceutically acceptable carrier.

The compounds according to the present invention may be used in several indications which involve 11- β -hydroxysteroid dehydrogenase type 1 enzyme. Thus the
15 compounds according to the present invention may be used against dementia (see WO97/07789), osteoporosis (see Canalis E 1996, Mechanisms of glucocorticoid action in bone: implications to glucocorticoid-induced osteoporosis, Journal of Clinical Endocrinology and Metabolism, 81, 3441-3447) and may also be used disorders in the immune system (see Franchimont et al, "Inhibition of Th1 immune response by
20 glucocorticoids: dexamethasone selectively inhibits IL-12-induced Stat 4 phosphorylation in T lymphocytes", The journal of Immunology 2000, Feb 15, vol 164 (4), pages 1768-74) and also in the above listed indications.

The various terms used, separately and in combinations, in the above definition of the
25 compounds having the formula (I) will be explained.

The term "aryl" in the present description is intended to include aromatic rings (monocyclic or bicyclic) having from 6 to 10 ring carbon atoms, such as phenyl (Ph) and naphthyl, which optionally may be substituted by C₁₋₆-alkyl. Examples of
30 substituted aryl groups are benzyl, and 2-methylphenyl.

The term "heteroaryl" means in the present description a monocyclic, bi- or tricyclic aromatic ring system (only one ring need to be aromatic) having from 5 to 14, preferably 5 to 10 ring atoms such as 5, 6, 7, 8, 9 or 10 ring atoms (mono- or bicyclic), in which one or more of the ring atoms are other than carbon, such as nitrogen, sulfur, oxygen and selenium. Examples of such heteroaryl rings are pyrrole, imidazole, thiophene, furan, thiazole, isothiazole, thiadiazole, oxazole, isoxazole, oxadiazole, pyridine, pyrazine, pyrimidine, pyridazine, pyrazole, triazole, tetrazole, chroman, isochroman, quinoline, quinoxaline, isoquinoline, phthalazine, cinnoline, quinazoline, indole, isoindole, indoline, isoindoline, benzothiophene, benzofuran, isobenzofuran, benzoxazole, 2,1,3-benzoxadiazole, benzothiazole, 2,1,3-benzothiazole, 2,1,3-benzoselenadiazole, benzimidazole, indazole, benzodioxane, indane, 1,2,3,4-tetrahydroquinoline, 3,4-dihydro-2*H*-1,4-benzoxazine, 1,5-naphthyridine, 1,8-naphthyridine, acridine, fenazine and xanthene.

The term "heterocyclic" in the present description is intended to include unsaturated as well as partially and fully saturated mono-, bi- and tricyclic rings having from 4 to 14, preferably 4 to 10 ring atoms, such as, for example, the heteroaryl groups mentioned above as well as the corresponding partially saturated or fully saturated heterocyclic rings. Exemplary saturated heterocyclic rings are azetidine, pyrrolidine, piperidine, piperazine, morpholine, thiomorpholine and 1,4-oxazepane.

C₁₋₆-alkyl in the compound of formula (I) according to the present application, which may be straight, branched or cyclic, is preferably C₁₋₄-alkyl. Exemplary alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, isopentyl, hexyl, isohexyl, and cyclohexyl.

C₁₋₆-alkoxy, in the compound of formula (I) according to the present application may be straight or branched, is preferably C₁₋₄-alkoxy. Exemplary alkoxy groups include methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, hexyloxy, and isohexyloxy.

C₁₋₆-acyl, in the compound of formula (I) according to the present application may be saturated or unsaturated and is preferably C₁₋₄-acyl. Exemplary acyl groups include formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, butenoyl (e.g. 3-butenoyl), hexenoyl (e.g. 5-hexenoyl).

5

The term "halogen" in the present description is intended to include fluorine, chlorine, bromine and iodine.

10 With the expression mono- or di-substituted is meant in the present description that the functionalities in question may be substituted with independently H, C₁₋₆-acyl, C₁₋₆-alkenyl, C₁₋₆-(cyclo)alkyl, aryl, pyridylmethyl, or heterocyclic rings e.g. azetidine, pyrrolidine, piperidine, piperazine, morpholine and thiomorpholine, which heterocyclic rings optionally may be substituted with C₁₋₆-alkyl.

15 The term "prodrug forms" in the present description means a pharmacologically acceptable derivative, such as an ester or an amide, which derivative is biotransformed in the body to form the active drug (see Goodman and Gilman's, The Pharmacological basis of Therapeutics, 8th ed., McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs, p. 13-15).

20

"Pharmaceutically acceptable" means in the present description being useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes being useful for veterinary use as well as human pharmaceutical use.

25

"Pharmaceutically acceptable salts" mean in the present description salts which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity. Such salts include acid addition salts formed with organic and inorganic acids, such as hydrogen chloride, hydrogen bromide, hydrogen iodide, 30 sulfuric acid, phosphoric acid, acetic acid, glycolic acid, maleic acid, malonic acid, oxalic acid, methanesulfonic acid, trifluoroacetic acid, fumaric acid, succinic acid,

tartaric acid, citric acid, benzoic acid, ascorbic acid and the like. Base addition salts may be formed with organic and inorganic bases, such as sodium, ammonia, potassium, calcium, ethanolamine, diethanolamine, N-methylglucamine, choline and the like.

5

Pharmaceutical compositions according to the present invention contain a pharmaceutically acceptable carrier together with at least one of the compounds comprising the formula (I) as described herein above, dissolved or dispersed therein as an active, antimicrobial, ingredient. In a preferred embodiment, the therapeutic composition is not immunogenic when administered to a human patient for therapeutic purposes, unless that purpose is to induce an immune response.

The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art. Typically such compositions are prepared as sterile injectables either as liquid solutions or suspensions, aqueous or non-aqueous, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be emulsified.

The active ingredient may be mixed with excipients, which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if desired, the composition may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like which enhance the effectiveness of the active ingredient. Adjuvants may also be present in the composition.

Pharmaceutically acceptable carriers are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at physiological

pH value, physiological saline or both, such as phosphate-buffered saline. Still further, aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and potassium chlorides, dextrose, propylene glycol, polyethylene glycol and other solutes.

5

Liquid compositions can also contain liquid phases in addition to and to the exclusion of water. Exemplary of such additional liquid phases are glycerine, vegetable oils such as cottonseed oil, organic esters such as ethyl oleate, and water-oil emulsions.

10 The pharmaceutical composition according to one of the preferred embodiments of the present invention comprising compounds comprising the formula (I), may include pharmaceutically acceptable salts of that component therein as set out above.

Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for
15 example, hydrochloric or phosphoric acids, or such organic acids as acetic acid, tartaric acid, mandelic acid and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

20

The preparations according to the preferred embodiments may be administered orally, topically, intraperitoneally, intraarticularly, intracranially, intradermally, intramuscularly, intraocularly, intrathecally, intravenously, subcutaneously. Other routes which are known for the skilled person in the art are thinkable.

25

The orally administrable compositions according to the present invention may be in the form of tablets, capsules, powders, granules, lozenges, liquid or gel preparations, such as oral, topical or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form and may contain

30 conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, traganath or polyvinyl-pyrrolidone; fillers e.g. lactose, sugar, maize-starch,

calcium phosphate, sorbitol or glycine; tableting lubricant e.g. magnesium stearate, talc, polyethylene glycol or silica; disintegrants e.g. potato starch, or acceptable wetting agents such as sodium lauryl sulfate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of e.g. aqueous or oily suspensions, solutions, emulsions, syrups or elixirs or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, e.g. sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible fats; emulsifying agents e.g. lecithin, sorbitan monooleate or acacia, non-aqueous vehicles (which may include edible oils), e.g. almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives e.g. methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

A pharmaceutical composition according to the present invention, may comprise typically an amount of at least 0.1 weight percent of compound comprising the formula (I) per weight of total therapeutic composition. A weight percent is a ratio by weight of total composition. Thus, for example, 0.1 weight percent is 0.1 grams of compound comprising the formula (I) per 100 grams of total composition. A suitable daily oral dose for a mammal, preferably a human being, may vary widely depending on the condition of the patient. However a dose of compound comprising the formula (I) of about 0.1 to 300 mg/kg body weight may be appropriate.

The compositions according to the present invention may also be used veterinarily and thus they may comprise a veterinarily acceptable excipient or carrier.

The compounds of the present invention in labelled form, e.g. isotopically labelled, may be used as a diagnostic agent.

The compounds of the formula (I) above may be prepared by, or in analogy with, conventional methods, and especially according to or in analogy with the following

methods. Further, the pharmacology in-vitro was studied using the following reagents and methods.

5 All publications mentioned herein are hereby incorporated by reference. By the expression "comprising" we understand including but not limited to. Thus, other non-mentioned substances, additives or carriers may be present.

10 The invention will now be described in reference to the following Figures and Examples. These Figures and Examples are not to be regarded as limiting the scope of the present invention, but shall only serve in an illustrative manner.

EXPERIMENTAL METHODS

Scintillation Proximity Assay

15 [1, 2(n) - ^3H]-cortisone was purchased from Amersham Pharmacia Biotech. Anti-cortisol monoclonal mouse antibody, clone 6D6.7 was obtained from Immunotech and Scintillation proximity assay (SPA) beads coated with monoclonal antimouse antibodies were from Amersham Pharmacia Biotech. NADPH, tetrasodium salt was
20 from Calbiochem and glucose-6-phosphate (G-6-P) was supplied by Sigma. The human 11- β -hydroxysteroid dehydrogenase type-1 enzyme (11- β -HSD₁) was expressed in *Pichia pastoris*. 18- β -glycyrrhetic acid (GA) was obtained from Sigma. The serial dilutions of the compounds were performed on a Tecan Genesis RSP 150. Compounds to be tested were dissolved in DMSO (1 mM) and diluted in 50 mM Tris-HCl, pH 7.2 containing 1 mM EDTA.
25

The multiplication of plates was done on a WallacQuadra. The amount of the product [^3H]-cortisol, bound to the beads was determined in a Packard, Top Count microplate liquid scintillation counter.

The 11- β -HSD₁ enzyme assay was carried out in 96 well microtiter plates (Packard, Optiplate) in a total well volume of 220 μ L and contained 30 mM Tris-HCl, pH 7.2 with 1 mM EDTA, a substrate mixture tritiated Cortisone/NADPH (175 nM / 181 μ M), G-6-P (1 mM) and inhibitors in serial dilutions (9 to 0.15 μ M). Reactions were initiated by the addition of human 11- β -HSD₁, either as *Pichia pastoris* cell homogenate or microsomes prepared from *Pichia pastoris* (the final amount of enzyme used was varied between 0.057 to 0.11 mg/mL). Following mixing, the plates were shaken for 30 to 45 minutes at room temperature. The reactions were terminated with 10 μ L 1 mM GA stop solution. Monoclonal mouse antibody was then added (10 μ L of 4 μ M) followed by 100 μ L of SPA beads (suspended according to the manufacturers instructions). Appropriate controls were set up by omitting the 11- β -HSD₁ to obtain the non-specific binding (NSB) value.

The plates were covered with plastic film and incubated on a shaker for 30 minutes, at room temperature, before counting. The amount of [³H]-cortisol, bound to the beads was determined in a microplate liquid scintillation counter.

The calculation of the K_i values for the inhibitors was performed by use of Activity Base. The K_i value is calculated from IC₅₀ and the K_m value is calculated using the Cheng Prushoff equation (with reversible inhibition that follows the Michaelis-Menten equation): $K_i = IC_{50}(1+[S]/K_m)$ [Cheng, Y.C.; Prushoff, W.H. Biochem. Pharmacol. 1973, 22, 3099-3108]. The IC₅₀ is measured experimentally in an assay wherein the decrease of the turnover of cortisone to cortisol is dependent on the inhibition potential of each substance. The K_i values of the compounds of the present invention for the 11- β -HSD₁ enzyme lie typically between about 10 nM and about 10 μ M. Illustrative of the invention, the following K_i values have been determined in the human 11- β -HSD₁ enzyme assay (see Table 1):

Table 1: K_i values determined in the human 11- β -HSD₁ enzyme assay.

Compound of Example	K _i (nM)
14	163
28	14
30	91

COMPOUND PREPARATION

General:

- 5 For preparative straight phase HPLC purification a Phenomenex column (250 × 21.1 mm, 10 μm) was used on a Gilson system eluting with ethanol in chloroform (gradient from 0 – 10% in 10 min) with a flow of 20 mL/min. Column chromatography was performed on silica using Silica gel 60 (230-400 mesh), Merck. Melting points were determined on a Gallenkamp apparatus. Elemental analyses were recorded using a
- 10 Vario EL instrument. HPLC analyses were performed using a Hypersil Elite column (150 x 4.6 mm, 3μ) with a flow of 3 mL / min on a Waters 600E system with monitoring at 254 nm. Reverse phase preparative HPLC was carried out on a 100 x 21.2 mm, 5μ Hypersil Elite column eluting with a gradient of 5% ACN in 95% water to 95% ACN in 5% water (0.2% TFA buffer) over 10 mins at a flow rate of 20 mL /
- 15 min with the UV detector set at 254 nm. Thin layer chromatography was carried out using pre-coated silica gel F-254 plates (thickness 0.25 mm). Electrospray MS spectra were obtained on a Micromass platform LCMS spectrometer. Crude, worked up compounds were purified by flash column chromatography using pre packed silica SPE columns (10 g silica) on an Isco Foxy 200 Combiflash system, and a gradient of
- 20 16.67% ethyl acetate in hexane increasing incrementally to 100% ethyl acetate.

List of Abbreviations

DCM = dichloromethane

25 DMAP = 4-dimethylaminopyridine

DMF = dimethylformamide

DMSO = dimethyl sulfoxide

EDTA = ethylenediaminetetraacetic acid

SULFONAMIDE COUPLINGS:

5 METHOD A:

1 Eq of the 2-aminothiazole was dissolved in pyridine (0.5 M solution). The sulfonyl chloride (1.2 eq) was added and the reaction mixture was stirred at ambient temperature under nitrogen atmosphere for 15 h. The reaction mixture was poured into aqueous HCl (1 M). If the product precipitated it was collected on a filter and washed
10 with aqueous HCl (1 M) and recrystallised from ethanol. In case an oil was obtained, the crude was extracted with DCM and worked up and purified using standard procedures.

METHOD B:

15 A solution of the 2-aminothiazole derivative (1 eq), triethylamine (2 eq) and DMAP (1 eq) in DMF (1 M) and DCM (0.225 M) was dispensed into a reaction vial. The sulfonyl chloride (1.2 eq) was dissolved in DCM (0.33 M) and added. The reaction mixtures were kept at room temperature over night. The mixture was then added to petroleum ether (10 times reaction volume). After some hours in refrigerator the
20 supernatants were decanted and (a portion of) the residual materials were dissolved in DMSO-methanol-acetic acid (300 μ L + 500 μ L + 50 μ L) and purified by preparative LCMS (acetonitrile-water gradients). The purest fractions were collected and lyophilized. Alternatively, the crude was isolated using extractive work up and purified using standard procedures.

25

FORMATION OF THIAZOLE RING:

METHOD H:

To a solution or suspension of a (substituted) thiourea in ethanol (0.5 M), 1 equivalent
30 of α -haloketone was added at room temperature. The reaction mixture was stirred in a sealed tube at 95 °C for 4 h, cooled, concentrated, redissolved in ethyl acetate, washed

with saturated aqueous sodium hydrogen carbonate, dried over sodium sulfate and chromatographed on silica gel using petroleum-ether and ethyl acetate as eluents.

METHOD I:

- 5 To a 0.5 M solution of ketone (1 eq) and thiourea (2 eq) in ethanol at 60 °C, 1 eq of iodine was added in one portion. The reaction tube was sealed and the reaction mixture was stirred at 100 °C for 16 hours. After evaporation of the solvent the residue was taken up in DCM, washed with saturated aqueous sodium hydrogen carbonate, dried with magnesium sulfate. Products were purified by chromatography on silica gel using
10 a gradient of petroleum-ether / ethyl acetate from 8:1 to 2:1 for elution.

KNOWN EXAMPLES

- The compounds of the following Examples 1-3 [244A-246A] are commercially
15 available and could e g be purchased from Bionet (Example 1 and 2) or Maybridge (Example 3).

- 1 [244A] Ethyl 2-(2-{[(4-chlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)benzoate
2 [245A] 2,5-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide
20 3 [246A] 4-Chloro-N-[4-(4,5-dichloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide

NOVEL EXAMPLES

- The following specific compounds were synthesized. The commercially available
25 compounds thus only form embodiments, as indicated earlier in the description, as pharmaceutical preparations and use of said compounds as set out in the appended set of claims.

EXAMPLE 4 [248A]

Ethyl 2-[[[(4-chlorophenyl)sulfonyl]amino]-4-phenyl-1,3-thiazole-5-carboxylate

The title compound was prepared from ethyl 2-amino-4-phenylthiazole-5-carboxylate and 4-chlorobenzenesulfonyl chloride as described in the synthetic METHOD B to
5 give a white solid (61.7 mg) with purity >90%. LCMS (pos) m/z 423.0.

EXAMPLE 5 [250A]

Ethyl 2-[[[(3-chloro-2-methylphenyl)sulfonyl]amino]-4-phenyl-1,3-thiazole-5-carboxylate

10 The title compound was prepared from ethyl 2-amino-4-phenylthiazole-5-carboxylate and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (47.2 mg) with purity >90%. LCMS (pos) m/z 437.0.

15 **EXAMPLE 6 [252A]**

N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 2-amino-4-(3-nitrophenyl)thiazole and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (41.5 mg) with purity >90%; MS (pos) m/z 404.1, (neg) m/z 402.2;
20 HRMS m/z 403.0062 (calc. of monoisotopic mass for C₁₈H₁₇N₃O₄S₂ gives 403.0060).

EXAMPLE 7 [253A]

N-(4-phenyl-1,3-thiazol-2-yl)-4-propylbenzenesulfonamide

The title compound was prepared from 2-amino-4-phenylthiazole and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
25 white solid (29.8 mg) with purity >90%. MS (pos) m/z 359.2.

EXAMPLE 8 [254A]

N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 2-amino-4-(4'-fluoro-3'-methyl)phenylthiazole and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to
5 give a white solid (31.5 mg) with purity >90%. MS (pos) m/z 391.2.

EXAMPLE 9 [255A]

N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 4-(4-methoxy-phenyl)-thiazol-2-ylamine and
10 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (32.2 mg) with purity >90%. MS (pos) m/z 389.2.

EXAMPLE 10 [256A]

3-Chloro-2-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide

15 The title compound was prepared from 2-amino-4-(-3-nitrophenyl)thiazole and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (36.0 mg) with purity >90%. MS (pos) m/z 410.0.

EXAMPLE 11 [257A]

20 **3-Chloro-2-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide**

The title compound was prepared from 2-amino-4-phenylthiazole and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (51.8 mg) with purity >90%. MS (pos) m/z 365.1.

25 **EXAMPLE 12 [258A]**

3-Chloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide

The title compound was prepared from 2-amino-4-(4'-fluoro-3'-methyl)phenylthiazole and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic
30 METHOD B to give a white solid (39.7 mg) with purity >90%. MS (pos) m/z 397.1.

EXAMPLE 13 [259A]**2,4,6-Trichloro-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide**

The title compound was prepared from 2-amino-4-(3-nitrophenyl)thiazole and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a
5 yellow solid (16.2 mg) with purity >90%. MS (pos) m/z 465.9.

EXAMPLE 14 [260A]**2,4,6-Trichloro-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide**

The title compound was prepared from 2-amino-4-phenylthiazole and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a
10 white solid (16.0 mg) with purity >90%. MS (pos) m/z 419.0, 421.0, 423.0.

EXAMPLE 15 [261A]**2,4,6-Trichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]benzenesulfonamide**
15

The title compound was prepared from 2-amino-4-(4'-fluoro-3'-methyl)phenylthiazole and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (50.9 mg) with purity >90%. MS (pos) m/z 451.0, 453.0, 455.0; HRMS m/z 449.9218 (calc. of monoisotopic mass for C₁₆H₁₀Cl₃FN₂O₂S₂ gives
20 449.9233).

EXAMPLE 16 [262A]**2,4,6-Trichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]benzenesulfonamide**

The title compound was prepared from 4-(4-methoxy-phenyl)-thiazol-2-ylamine and
25 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a white-yellow solid (58.2 mg) with purity >90%. MS (pos) m/z 449.1, 451.1, 453.1.

EXAMPLE 17 [263A]**N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide**

The title compound was prepared from 2-amino-4-(-3-nitrophenyl)thiazole and 4-phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
5 yellow solid (25.8 mg) with purity >80%. MS (pos) m/z 438.1.

EXAMPLE 18 [264A]**N-(4-phenyl-1,3-thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide**

The title compound was prepared from 2-amino-4-phenylthiazole and 4-
10 phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (32.9 mg) with purity >90%. MS (pos) m/z 393.2; HRMS m/z 392.0658 (calc. of monoisotopic mass for C₂₁H₁₆N₂O₂S₂ gives 392.0653).

EXAMPLE 19 [265A]**15 N-[4-(4-Fluoro-3-methylphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide**

The title compound was prepared from 2-amino-4-(4'-fluoro-3'-methyl)phenylthiazole and 4-phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (15.8 mg) with purity >90%. MS (pos) m/z 425.2.

20 EXAMPLE 20 [266A]**N-[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide**

The title compound was prepared from 4-(4-methoxy-phenyl)-thiazol-2-ylamine 4-phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (17.8 mg) with purity >90%. MS (pos) m/z 423.3.

25

EXAMPLE 21 [267A]**2,4-Dichloro-6-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide**

The title compound was prepared from 2-amino-4-(-3-nitrophenyl)thiazole and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic METHOD B
30 to give a yellow solid (27.3 mg) with purity >90%. MS (pos) m/z 444.0, 446.0.

EXAMPLE 22 [268A]**2,4-Dichloro-6-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide**

The title compound was prepared from 2-amino-4-phenylthiazole and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
5 white solid (49.2 mg) with purity >90%. MS (pos) m/z 399.1, 401.1.

EXAMPLE 23 [269A]**2,4-Dichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide**

10 The title compound was prepared from 2-amino-4-(4'-fluoro-3'-methyl)phenylthiazole and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (28.4 mg) with purity >90%. MS (pos) m/z 431.1, 433.1.

15 EXAMPLE 24 [270A]**2,4-Dichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide**

The title compound was prepared from 4-(4-methoxy-phenyl)-thiazol-2-ylamine 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic METHOD B
20 to give a white solid (48.4 mg) with purity >90%. MS (pos) m/z 429.1, 431.1; HRMS m/z 427.9810 (calc. of monoisotopic mass for $C_{17}H_{14}Cl_2N_2O_3S_2$ gives 427.9823).

EXAMPLE 25 [271A]**N-[4-(2-[(4-propylphenyl)sulfonyl]amino)-1,3-thiazol-4-yl]phenyl]acetamide**

25 The title compound was prepared from N-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (1.0 mg) with purity >80%. MS (pos) m/z 416.2.

EXAMPLE 26 [272A]**4-Propyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide**

The title compound was prepared from 4-(3-pyridyl)-1,3-thiazol-2-amine and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
5 white-yellow solid (2.3 mg) with purity >90%. MS (pos) m/z 360.2.

EXAMPLE 27 [273A]**N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide**

The title compound was prepared from 4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-amine
10 hydrobromide and 4-n-propylbenzenesulfonyl chloride as described in the synthetic
METHOD B to give a white solid (9.0 mg) with purity >90%. MS (pos) m/z 438.2.

EXAMPLE 28 [274A]**N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-
15 propylbenzenesulfonamide**

The title compound was prepared from 7-methoxy-4,5-dihydronaphtho[1,2-
D][1,3]thiazol-2-amine hydrobromide and 4-n-propylbenzenesulfonyl chloride as
described in the synthetic METHOD B to give a yellow solid (29.0 mg) with purity
>90%: MS (pos) m/z 415.3; HRMS m/z 414.1068 (calc. of monoisotopic mass for
20 $C_{21}H_{22}N_2O_3S_2$ gives 414.1072).

EXAMPLE 29 [275A]**N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide**

The title compound was prepared from 4-(5-chloro-2-thienyl)-1,3-thiazol-2-amine and
25 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give
a brown solid (16.4 mg) with purity >80%. MS (pos) m/z 399.2.

EXAMPLE 30 [276A]

N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
5 white solid (30.9 mg) with purity >90%. MS (pos) m/z 393.1.

EXAMPLE 31 [277A]

N-[4-(2-[(3-chloro-2-methylphenyl)sulfonyl]amino)-1,3-thiazol-4-yl)phenyl]acetamide

10 The title compound was prepared *N*-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (16.1 mg) with purity >90%. MS (pos) m/z 422.1, (neg) m/z 420.3. HRMS m/z 421.0327 (calc. of monoisotopic mass for $C_{18}H_{16}ClN_3O_3S_2$ gives 421.0322).

15

EXAMPLE 32 [278A]

3-Chloro-2-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(3-pyridyl)-1,3-thiazol-2-amine and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to
20 give a yellow solid (21.4 mg) with purity >90%. MS (pos) m/z 366.1.

EXAMPLE 33 [279A]

3-Chloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide

25 The title compound was prepared from 4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-amine hydrobromide and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (16.5 mg) with purity >90%. 1H -NMR (DMSO- d_6) δ 13.43 (s, NH), 8.46 (d, $J=2.8$ Hz, 1H), 8.29 (d, $J=2.8$ Hz, 1H), 7.96 (dd, $J=1.1$ Hz, $J=8.0$ Hz, 1H), 7.90 (d, $J=8.8$ Hz, 1H), 7.70 (dd, $J=1.1$ Hz, $J=8.0$ Hz, 1H),
30 7.42 (t, $J=8.0$ Hz, 1H), 7.52 (s, 1H), 2.67 (s, 3H); MS (pos) m/z 442.2, 444.2.

EXAMPLE 34 [280A]

3-Chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-methylbenzenesulfonamide

The title compound was prepared from 7-methoxy-4,5-dihydronaphtho[1,2-D][1,3]thiazol-2-amine hydrobromide and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (30.7 mg) with purity >90%. MS (pos) m/z 421.1.

EXAMPLE 35 [281A]

3-Chloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide

The title compound was prepared from 4-(5-chloro-2-thienyl)-1,3-thiazol-2-amine and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a beige solid (29.9 mg) with purity >90%: MS (pos) m/z 405.0, 407.0; HRMS m/z 403.9278 (calc. of monoisotopic mass for C₁₄H₁₀Cl₂N₂O₂S₃ gives 403.9281).

EXAMPLE 36 [282A]

3-Chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide

The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (45.3 mg) with purity >90%: MS (pos) m/z 399.1, 401.1; HRMS m/z 397.9711 (calc. of monoisotopic mass for C₁₆H₁₂Cl₂N₂O₂S₂ gives 397.9117).

EXAMPLE 37 [283A]

N-[4-(2-[(2,4,6-trichlorophenyl)sulfonyl]amino)-1,3-thiazol-4-yl)phenyl]acetamide

The title compound was prepared from N-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (2.5 mg) with purity >80%. MS (pos) m/z 476.1, 478.1, 480.1.

EXAMPLE 38 [284A]

2,4,6-Trichloro-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(3-pyridyl)-1,3-thiazol-2-amine and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a
5 yellow solid (29.3 mg) with purity >90%. MS (pos) m/z 422.1; HRMS m/z 418.9134 (calc. of monoisotopic mass for C₁₄H₈Cl₃N₃O₂S₂ gives 418.9123).

EXAMPLE 39 [285A]

2,4,6-Trichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-
10 yl]benzenesulfonamide

The title compound was prepared from 4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-amine hydrobromide and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (11.3 mg) with purity >90%. MS (pos)
m/z 498.0, 500.0, 502.0.

15

EXAMPLE 40 [286A]

2,4,6-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-
yl)benzenesulfonamide

The title compound was prepared from 7-methoxy-4,5-dihydronaphtho[1,2-
20 D][1,3]thiazol-2-amine hydrobromide and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (10.1 mg) with purity >90%. MS (pos) m/z 477.1.

EXAMPLE 41 [287A]

25 2,4,6-Trichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(5-chloro-2-thienyl)-1,3-thiazol-2-amine and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a yellow solid (24.0 mg) with purity >90%. MS (pos) m/z 461.0, 463.0.

EXAMPLE 42 [289A]

2,4,6-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine and 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to
5 give a white solid (38.7 mg) with purity >90%. MS (pos) m/z 453.0, 455.0, 457.0.

EXAMPLE 43 [290A]

N-(4-{2-[(1,1'-biphenyl)-4-ylsulfonyl]amino}-1,3-thiazol-4-yl)phenylacetamide

The title compound was prepared *N*-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide
10 and 4-phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to
give a white-yellow solid (8.6 mg) with purity >80%. MS (pos) m/z 450.3.

EXAMPLE 44 [291A]

N-[4-(3-pyridinyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide

15 The title compound was prepared from 4-(3-pyridyl)-1,3-thiazol-2-amine and 4-phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
white-yellow solid (1.4 mg) with purity >80%. MS (pos) m/z 394.2.

EXAMPLE 45 [292A]

20 N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide

The title compound was prepared from 4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-amine hydrobromide and 4-phenylbenzenesulfonyl chloride as described in the synthetic
METHOD B to give a white solid (11.9 mg) with purity >90%. MS (pos) m/z 472.1.

25 EXAMPLE 46 [293A]

N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide

The title compound was prepared from 7-methoxy-4,5-dihydronaphtho[1,2-D][1,3]thiazol-2-amine hydrobromide and 4-phenylbenzenesulfonyl chloride as
30 described in the synthetic METHOD B to give a white-yellow solid (17.0 mg) with
purity >80%. MS (pos) m/z 449.2.

EXAMPLE 47 [294A]**N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide**

The title compound was prepared from 4-(5-chloro-2-thienyl)-1,3-thiazol-2-amine and
5 4-phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
white solid (4.3 mg) with purity >80%. MS (pos) m/z 433.1, 435.1.

EXAMPLE 48 [295A]**N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide**

10 The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine and 4-
phenylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a
white solid (31.8 mg) with purity >90%. MS (pos) m/z 427.1.

EXAMPLE 49 [296A]**N-[4-(2-((2,4-dichloro-6-methylphenyl)sulfonyl)amino)-1,3-thiazol-4-
15 yl)phenyl]acetamide**

The title compound was prepared N-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide
and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic
METHOD B to give a white-yellow solid (8.1 mg) with purity >80%. MS (pos) m/z
20 456.2, 458.1.

EXAMPLE 50 [297A]**2,4-Dichloro-6-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide**

The title compound was prepared from 4-(3-pyridyl)-1,3-thiazol-2-amine and 2,4-
25 dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic METHOD B
to give a yellow solid (17.6 mg) with purity >90%. MS (pos) m/z 400.0, 402.0.

EXAMPLE 51 [298A]

2,4-Dichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide

The title compound was prepared from 4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-amine
5 hydrobromide and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the
synthetic METHOD B to give a white-yellow solid (18.4 mg) with purity >90%. MS
(pos) m/z 478.0, 479.9.

EXAMPLE 52 [299A]

10 **2,4-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-6-methylbenzenesulfonamide**

The title compound was prepared from 7-methoxy-4,5-dihydronaphtho[1,2-
D][1,3]thiazol-2-amine hydrobromide and 2,4-dichloro-6-methylbenzenesulfonyl
chloride as described in the synthetic METHOD B to give a white solid (16.3 mg) with
15 purity >90%. MS (pos) m/z 455.1, 457.1.

EXAMPLE 53 [300A]

2,4-Dichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide

20 The title compound was prepared from 4-(5-chloro-2-thienyl)-1,3-thiazol-2-amine
(METHOD H) and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the
synthetic METHOD B to give a white solid (19.5 mg) with purity >90%. MS (pos)
m/z 439.0, 441.0.

25 EXAMPLE 54 [301A]

2,4-Dichloro-N-[4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide

The title compound was prepared from 4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-ylamine
(METHOD I) and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the
30 synthetic METHOD B to give a yellow solid (11.6 mg) with purity >90%. MS (pos)
m/z 417.1, 419.1.

EXAMPLE 55 [304A]

N-[4-(1-benzothien-3-yl)-1,3-thiazol-2-yl]-2,4-dichloro-6-methylbenzenesulfonamide

- 5 The title compound was prepared from 4-(1-benzothien-3-yl)-1,3-thiazol-2-amine (METHOD I) and 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (47.0 mg) with purity >90%. MS (pos) m/z 455.0, 457.0.

10 **EXAMPLE 56 [305A]**

N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine and 4-n-propylbenzenesulfonyl chloride as described in the synthetic METHOD B to give a brown-red solid (5.3 mg) with purity >90%. MS (pos) m/z 399.2, 401.2.

15

EXAMPLE 57 [306A]

3-Chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide

- The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine and 3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B
20 to give a white-yellow solid (7.6 mg) with purity >90%. MS (pos) m/z 405.2, 407.2.

EXAMPLE 58 [307A]

2,4,6-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide

- The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine and
25 2,4,6-trichlorobenzenesulfonyl chloride as described in the synthetic METHOD B to give a white solid (5.0 mg) with purity >90%. MS (pos) m/z 459.1, 461.1, 463.1.

EXAMPLE 59 [308A]**2,4-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide**

The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine and
5 2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic
METHOD B to give a yellow solid (9.4 mg) with purity >90%. MS (pos) m/z 439.1,
441.1; HRMS m/z 437.8875 (calc. of monoisotopic mass for C₁₄H₉Cl₃N₂O₂S₃ gives
437.8892).

10 EXAMPLE 60 [311A]**2,4-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide**

The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine and
2,4-dichloro-6-methylbenzenesulfonyl chloride as described in the synthetic
METHOD B to give a white solid (40.3 mg) with purity >90%. MS (pos) m/z 435.0,
15 437.0.

EXAMPLE 61 [312A]**Ethyl 2-([(1,1'-biphenyl)-4-ylsulfonyl]amino)-4-phenyl-1,3-thiazole-5-carboxylate**

The title compound was prepared from 2-amino-4-phenylthiazole-5-carboxylate and 4-
20 biphenylsulfonyl chloride as described in the synthetic METHOD B to give a white
solid (21.7 mg) with purity >90%. LCMS (pos) m/z 465.2.

EXAMPLE 62 [313A]**3-Chloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide**

25 The title compound was prepared from 4-(4-methoxy-phenyl)-thiazol-2-ylamine and
3-chloro-2-methylbenzenesulfonyl chloride as described in the synthetic METHOD B
to give a white solid (58.0 mg) with purity >90%. MS (pos) m/z 395.2.

EXAMPLE 63 [313B]

N-[4-(2-[(4-Bromo-2,5-difluorophenyl)sulfonyl]amino)-1,3-thiazol-4-yl)phenyl]acetamide

The title compound was prepared *N*-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide (63 mg) and 4-bromo-2,5-difluorobenzenesulfonyl chloride (79 mg) as described in the synthetic METHOD B to give a white-yellow solid (6.2 mg) with purity >90%: MS (pos) *m/z* 488.3, 490.3.

EXAMPLE 64 [313C]

2,3,4-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine (57 mg) and 2,3,4-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (64.8 mg) with purity >90%: MS (pos) *m/z* 453.3, 455.3, 457.2.

EXAMPLE 65 [313D]

2,4,5-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine (59 mg) and 2,4,5-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (9.2 mg) with purity >90%: MS (pos) *m/z* 459.2, 461.2, 463.2.

EXAMPLE 66 [313E]

2,3,4-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine (59 mg) and 2,3,4-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (5.0 mg) with purity >90%: MS (pos) *m/z* 459.2, 461.2, 463.2.

EXAMPLE 67 [313F]

4-Bromo-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2,5-difluorobenzenesulfonamide

The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine (59 mg) and 4-bromo-2,5-difluorobenzenesulfonyl chloride (79 mg) as described in the synthetic METHOD B to give a white solid (22.3 mg) with purity >90%: MS (pos) m/z 471.2, 473.2.

EXAMPLE 68 [313G]

4,5-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide

The title compound was prepared from 7-methoxy-4,5-dihydronaphtho(1,2-D)(1,3)thiazol-2-amine hydrobromide (85 mg) and 2,3-dichlorothiophene-5-sulfonyl chloride (68 mg) as described in the synthetic METHOD B to give a white solid (16.4 mg) with purity >90%: MS (pos) m/z 447.3, 449.3.

EXAMPLE 69 [313H]

4,5-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide

The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine (57 mg) and 2,3-dichlorothiophene-5-sulfonyl chloride (68 mg) as described in the synthetic METHOD B to give a white solid (56.6 mg) with purity >90%: MS (pos) m/z 425.3, 427.3; HRMS m/z 423.8737 (calc. of monoisotopic mass for C₁₃ H₇ Cl₃ N₂ O₂ S₃ gives 423.8735).

25 EXAMPLE 70 [313I]

N-[4-(2-((2,4,5-Trichlorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl)phenyl]acetamide

The title compound was prepared from N-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide (63 mg) and 2,4,5-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white-yellow solid (5.0 mg) with purity >90%: MS (pos) m/z 476.3, 478.3.

EXAMPLE 71 [313J]

2,3,4-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide

- 5 The title compound was prepared from 7-methoxy-4,5-dihydronaphtho(1,2-D)(1,3)thiazol-2-amine hydrobromide (85 mg) and 2,3,4-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (13.1 mg) with purity >90%: MS (pos) m/z 475.3, 477.3

10 **EXAMPLE 72 [313K]**

4-Bromo-5-chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide

- The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine (59 mg) and 4-bromo-5-chlorothiophene-2-sulfonyl chloride (80 mg) as described in the
15 synthetic METHOD B to give a white solid (34.2 mg) with purity >90%: MS (pos) m/z 457.1, 477.1, 479.1.

EXAMPLE 73 [313L]

- 3-Bromo-5-chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide**
20

The title compound was prepared 7-methoxy-4,5-dihydronaphtho(1,2-D)(1,3)thiazol-2-amine hydrobromide (85 mg) and 3-bromo-5-chlorothiophene-2-sulfonyl chloride (80 mg) as described in the synthetic METHOD B to give a white solid (17.8 mg) with
25 purity >80%: MS (pos) m/z 491.3, 493.3.

EXAMPLE 74 [313M]

3-Bromo-5-chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide

- The title compound was prepared from 4-(2-chlorophenyl)-1,3-thiazol-2-amine (57
30 mg) and 3-bromo-5-chlorothiophene-2-sulfonyl chloride (80 mg) as described in the

synthetic METHOD B to give a white solid (43.7 mg) with purity >90%: MS (pos) m/z 469.2, 471.2, 473.2.

EXAMPLE 75 [313N]

5 N-[4-(2-{{(2,6-Dichlorophenyl)sulfonyl}amino}-1,3-thiazol-4-yl)phenyl]acetamide

The title compound was prepared from N-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide (63 mg) and 2,6-dichlorobenzenesulfonyl chloride (66 mg) as described in the synthetic METHOD B to give a white solid (5.6 mg) with purity >80%: MS (pos) m/z 442.2, 444.4.

10

EXAMPLE 76 [313O]

2,6-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide

The title compound was prepared from 4-(3-chloro-2-thienyl)-1,3-thiazol-2-amine (59 mg) and 2,6-dichlorobenzenesulfonyl chloride (66 mg) as described in the synthetic

15 METHOD B to give a yellow solid (34.5 mg) with purity >90%: MS (pos) m/z 425.3, 427.3; HRMS m/z 423.8730 (calc. of monoisotopic mass for C₁₃ H₇ Cl₃ N₂ O₂ S₃ gives 423.8735).

EXAMPLE 77 [313P]

20 2,4,6-Trichloro-N-(7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide

The title compound was prepared from 7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-amine (100 mg, synthesized according to METHOD I from 6,7-dimethoxytetralone) and 2,4,6-trichlorobenzenesulfonyl chloride (213 mg) as

25 described in the synthetic METHOD A to give a yellow powder (6.5 mg) with a purity of 86%: MS-ES (neg) m/z 505.1

EXAMPLE 78 [313Q]

2,3,4-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide

The title compound was prepared from 4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-amine (91 mg) and 2,3,4-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (49.2 mg) with purity >90%: MS (pos) m/z 579.1, 581.1, 583.1; MS (neg) m/z 577.5, 579.5, 581.5.

EXAMPLE 79 [313R]

2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide

The title compound was prepared from 4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-amine (85 mg) and 2,3,4-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (94.0 mg) with purity >90%: MS (pos) m/z 555.1, 557.1, 559.1; MS (neg) m/z 553.4, 555.4, 557.4; HRMS m/z 553.8262 (calc. of monoisotopic mass for C₁₆ H₆ Cl₅ F₃ N₂ O₂ S₂ gives 553.8265).

EXAMPLE 80 [313S]

N-[4-(2-Chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-amine (62 mg) and 4-n-propylbenzenesulfonyl chloride (59 mg) as described in the synthetic METHOD B to give a white solid (30.7 mg) with purity >90%: MS (pos) m/z 411.3, 413.3; MS (neg) m/z 409.5, 411.5.

EXAMPLE 81 [313T]

4-Bromo-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide

The title compound was prepared from 4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-amine (91 mg) and 4-bromo-2,5-difluorobenzenesulfonyl chloride (79 mg) as described in the synthetic METHOD B to give a white solid (46.9 mg) with purity

>90%: MS (neg) m/z 589.5, 591.5, 593.5; HRMS m/z 589.8745 (calc. of monoisotopic mass for C₂₁ H₁₁ Br Cl₂ F₂ N₂ O₃ S₂ gives 589.8740).

EXAMPLE 82 [313U]

5 4-Bromo-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide

The title compound was prepared 4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-amine (85 mg) and 4-bromo-2,5-difluorobenzenesulfonyl chloride (79 mg) as described in the synthetic METHOD B to give a white solid (38.2 mg) with purity

10 >90%: MS (pos) m/z 567.2, 569.2, 571.2; MS (neg) m/z 565.5, 567.5, 569.5.

EXAMPLE 83 [313V]

4,5-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide

15 The title compound was 4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-amine (62 mg) and 2,3-dichlorothiophene-5-sulfonyl chloride (68 mg) as described in the synthetic METHOD B to give a white solid (46.6 mg) with purity >90%: MS (pos) m/z 443.2, 445.2, 447.2; MS (neg) m/z 441.2, 443.2, 445.2; HRMS m/z 441.8657 (calc. of monoisotopic mass for C₁₃ H₆ Cl₃ F N₂ O₂ S₃ gives 441.8641).

20

EXAMPLE 84 [313W]

4-Bromo-5-chloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide

25 The title compound 4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-amine (91 mg) and 4-bromo-5-chlorothiophene-2-sulfonyl chloride (80 mg) as described in the synthetic METHOD B to give a white solid (44.4 mg) with purity >90%: MS (pos) m/z 595.3, 597.3, 599.3; MS (neg) m/z 593.2, 595.2, 597.2.

EXAMPLE 85 [313X]

4-Bromo-5-chloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide

The title compound 4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-amine
5 (85 mg) and 4-bromo-5-chlorothiophene-2-sulfonyl chloride (80 mg) as described in
the synthetic METHOD B to give a white solid (65.3 mg) with purity >90%: MS (pos)
m/z 571.2, 573.2, 575.2; MS (neg) m/z 569.1, 571.1, 573.3; HRMS m/z 569.7690
(calc. of monoisotopic mass for C₁₄ H₅ Br Cl₃ F₃ N₂ O₂ S₃ gives 569.7714).

10 EXAMPLE 86 [313Y]

2,4-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide

The title compound was 4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-amine (62 mg) and
2,4-dichloro-6-methylbenzenesulfonyl chloride (70 mg) as described in the synthetic
15 METHOD B to give a white solid (41.6 mg) with purity >90%: MS (pos) m/z 451.3,
453.3, 455.3; MS (neg) m/z 449.3, 451.3, 453.3.

EXAMPLE 87 [313Z]

20 **2,4,6-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide**

The title compound was 4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-
amine (85 mg) and 2,4,6-trichlorobenzenesulfonyl chloride (76 mg) as described in the
synthetic METHOD B to give a white solid (44.1 mg) with purity >90%: MS (pos)
m/z 555.3, 557.3, 559.3; MS (neg) m/z 553.2, 555.2, 557.2.

25

EXAMPLE 88 [313ZA]

4-Bromo-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide

The title compound was 4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-amine (62 mg)
30 and 4-bromo-2-methylbenzenesulfonyl chloride (73 mg) as described in the synthetic

METHOD B to give a white solid (18.0 mg) with purity >90%: MS (pos) m/z 461.3, 463.3, 465.4; MS (neg) m/z 459.3, 461.3, 463.3.

EXAMPLE 89 [313ZB]

5 2,4,6-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide

The title compound was 4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-amine (91 mg) and 2,4,6-trichlorobenzenesulfonyl chloride (76 mg) as described in the synthetic METHOD B to give a white solid (18.8 mg) with purity >90%: MS (pos)
10 m/z 578.9, 580.9, 582.9.

EXAMPLE 90 [313ZC]

N-[4-(2-{{(4-Bromo-5-chloro-2-thienyl)sulfonyl}amino}-1,3-thiazol-4-yl)phenyl]acetamide

15 The title compound was prepared *N*-[4-(2-amino-1,3-thiazol-4-yl)phenyl]acetamide (63 mg) and 4-bromo-5-chlorothiophene-2-sulfonyl chloride (80 mg) as described in the synthetic METHOD B to give a white solid (11.7 mg) with purity >80%: MS (pos) m/z 491.9, 493.9.

20 EXAMPLE 91 [313ZD]

N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide

The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (106 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-tetrahydrothianaphthene) and benzenesulfonyl chloride (93 mg) as described in the
25 synthetic METHOD A to give a white solid (13 mg) with a purity > 90%: MS-ES (neg) m/z 347.2.

EXAMPLE 92 [313ZE]

3,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide

The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (107 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-tetrahydrothianaphthene) and 3,4-dichlorobenzenesulfonyl chloride (127 mg) as described in the synthetic METHOD A to give a white powder (62 mg) with a purity > 90%: MS-ES (neg) m/z 417.2.

10 **EXAMPLE 93 [313ZF]**

3-Chloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-2-methylbenzenesulfonamide

The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (106 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-tetrahydrothianaphthene) and 3-chloro-2-methylbenzenesulfonyl chloride (114 mg) as described in the synthetic METHOD A to give a white powder (36 mg) with a purity > 90%: MS-ES (neg) m/z 395.2.

EXAMPLE 94 [313ZG]

20 **2,4,6-Trichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide**

The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (106 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-tetrahydrothianaphthene) and 2,4,6-trichlorobenzenesulfonyl chloride (141 mg) as described in the synthetic METHOD A to give a white powder (12 mg) with a purity > 90%: MS-ES (neg) m/z 451.2.

EXAMPLE 95 [313ZH]

N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide

30 The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (109 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-

tetrahydrothianaphthene) and 4-biphenylsulfonyl chloride (130 mg) as described in the synthetic METHOD A to give a white powder (12 mg) with a purity > 90%: MS-ES (neg) m/z 423.3.

5 **EXAMPLE 96 [313ZI]**

2,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-6-methylbenzenesulfonamide

The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (105 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-tetrahydrothianaphthene) and 2,4-dichloro-6-methylbenzenesulfonyl chloride (132 mg) as described in the synthetic METHOD A to give a white powder (21 mg) with a purity > 90%: MS-ES (neg) m/z 431.2.

EXAMPLE 97 [313ZJ]

15 **N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-4-propylbenzenesulfonamide**

The title compound was prepared from 4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-amine (105 mg, synthesized according to METHOD I from 4-keto-4,5,6,7-tetrahydrothianaphthene) and 4-n-propylbenzenesulfonyl chloride (111 mg) as described in the synthetic METHOD A to give a white powder (18 mg) with a purity > 90%: MS-ES (neg) m/z 389.3.

EXAMPLE 98 [313ZK]

3-Chloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-2-methylbenzenesulfonamide

25 The title compound was prepared from 6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-amine (61 mg, synthesized according to METHOD I from 1-chloro-3-(methylsulfonyl)-4,5,6,7-tetrahydrobenzo-[c]thiophen-4-one) and 3-chloro-2-methylbenzenesulfonyl chloride (47 mg) as described in the synthetic METHOD A to give a beige solid (24 mg) with a purity > 90%: MS-ES
30 (pos) m/z 509.1.

EXAMPLE 99 [313ZL]

N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-4-propylbenzenesulfonamide

The title compound was prepared from 6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-amine (61 mg, synthesized according to METHOD I from 1-chloro-3-(methylsulfonyl)-4,5,6,7-tetrahydrobenzo-[c]thiophen-4-one) and 4-n-propylbenzenesulfonyl chloride (46 mg) as described in the synthetic METHOD A to give a white powder (18 mg) with a purity of 90%: MS-ES (pos) m/z 501.2.

10

EXAMPLE 100 [313ZM]

2,4-Dichloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-6-methylbenzenesulfonamide

The title compound was prepared from 6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-amine (61 mg, synthesized according to METHOD I from 1-chloro-3-(methylsulfonyl)-4,5,6,7-tetrahydrobenzo-[c]thiophen-4-one) and 2,4-dichloro-6-methylbenzenesulfonyl chloride (54 mg) as described in the synthetic METHOD A to give a white solid (5.9 mg) with a purity > 90%: MS-ES (pos) m/z 543.1.

20

EXAMPLE 101 [313ZN]

N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl][1,1'-biphenyl]-4-sulfonamide

The title compound was prepared from 6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-amine (61 mg, synthesized according to METHOD I from 1-chloro-3-(methylsulfonyl)-4,5,6,7-tetrahydrobenzo-[c]thiophen-4-one) and 4-biphenylsulfonyl chloride (52 mg) as described in the synthetic METHOD A to give a beige solid (5.9 mg) with a purity > 90%: MS-ES (neg) m/z 535.2.

25

EXAMPLE 102 [313ZO]

N-{2-[(Phenylsulfonyl)amino]-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide

The title compound was prepared from 2-amino-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide (100 mg, synthesized according to METHOD I from 6-acetamidotetralone) and benzenesulfonyl chloride (81 mg) as described in the synthetic METHOD A to give a cream powder (4.8 mg) with a purity > 90%: MS-ES (neg) m/z 398.4.

10 EXAMPLE 103 [313ZP]

N-(2-[(3-Chloro-2-methylphenyl)sulfonyl]amino)naphtho[1,2-d][1,3]thiazol-6-yl}acetamide

The title compound was prepared from 2-amino-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide (100 mg, synthesized according to METHOD I from 6-acetamidotetralone) and 3-chloro-2-methylbenzenesulfonyl chloride (129 mg) as described in the synthetic METHOD A to give a cream powder (1.1 mg) with a purity of 87%: MS-ES (neg) m/z 444.3.

EXAMPLE 104 [313ZQ]

20 N-(2-[(4-Propylphenyl)sulfonyl]amino)-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide

The title compound was prepared from 2-amino-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide (100 mg, synthesized according to METHOD I from 6-acetamidotetralone) and 4-n-propylbenzenesulfonyl chloride (101 mg) as described in the synthetic METHOD A to give a cream powder (18 mg) with a purity > 90%: MS-ES (neg) m/z 440.4.

EXAMPLE 105 [313ZR]

N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide

The title compound was prepared from 8-nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-amine (100 mg, synthesized according to METHOD I from 7-nitrotetralone) and 4-n-propylbenzenesulfonyl chloride (106 mg) as described in the synthetic METHOD A to give a yellow powder (3.5 mg) with a purity of 87%: MS-ES (neg) m/z 428.4.

EXAMPLE 106 [313ZS]

N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide

The title compound was prepared from 8-nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-amine (100 mg, synthesized according to METHOD I from 7-nitrotetralone) and benzenesulfonyl chloride (85 mg) as described in the synthetic METHOD A to give a yellow powder (12 mg) with a purity of > 90%: MS-ES (neg) m/z 386.3.

EXAMPLE 107 [313ZT]


N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide

The title compound was prepared from 8-nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-amine (100 mg, synthesized according to METHOD I from 7-nitrotetralone) and 4-biphenylsulfonyl chloride (122 mg) as described in the synthetic METHOD A to give a yellow powder (4.4 mg) with a purity of > 90%: MS-ES (neg) m/z 462.4.

Various embodiments of the present invention have been described above but a person skilled in the art realizes further minor alterations which would fall into the scope of the present invention. The breadth and scope of the present invention should not be limited by any of the above-described exemplary embodiments, but should be defined only in accordance with the following claims and their equivalents.

Claims

1. A compound of the formula (I):



5

wherein

T is selected from

thienyl substituted with one or more of bromo, chloro;

10 phenyl substituted as follows:

- a) either T is phenyl, wherein the phenyl is substituted with one or more of propyl and phenyl;
- b) T is phenyl substituted with chloro in position 3 and methyl in position 2;
- 15 c) T is phenyl substituted with chloro in position 2 and 4, and methyl in position 6;
- d) T is phenyl substituted with bromo in position 4 and fluoro in position 2 and 5;
- e) T is phenyl substituted with chloro in position 2, 3, and 4;
- f) T is phenyl substituted with chloro in position 2, 4, and 5;
- g) T is phenyl substituted with bromo in position 4 and methyl in position 2;
- 20 h) T is phenyl substituted with chloro in position 2 and 6;
- i) T is phenyl substituted with chloro in position 2, 4, and 6; or
- j) T is phenyl substituted with bromo in position 4 and chloro in position 5.

A is selected from an aryl ring or heteroaryl ring, which can further be optionally
 25 substituted in one or more positions independently of each other by hydrogen, C₁₋₆-alkyl, halogenated C₁₋₆-alkyl, halogen, C₁₋₆-alkoxy, nitro, C₁₋₆-alkoxycarbonyl, C₁₋₆-alkylsulfonyl, acetylamino or aryloxy, wherein the aryloxy can further be optionally

substituted in one or more positions independently of each other by hydrogen and halogen;

B is selected from hydrogen and C₁₋₆-alkoxycarbonyl or is linked to A to give a 6-membered aromatic or non-aromatic ring;

as well as pharmaceutically acceptable salts, hydrates and solvates thereof.

2. A compound according to claim 1, wherein

A is selected from 1-benzothien-3-yl, 3-(2,5-dimethylfuryl), pyridinyl; thienyl optionally substituted with one or more of chloro, methylsulfonyl; phenyl optionally substituted with one or more of ethoxycarbonyl, nitro, fluoro, methyl, methoxy, acetylamino, chloro, 4-chlorophenoxy, trifluoromethyl;

B is selected from hydrogen, carbethoxy or is linked to A to give a 6-membered aromatic or non-aromatic ring.

3. A compound according to claim 1-2, selected from the group consisting of:

Ethyl 2-{[(3-chloro-2-methylphenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-carboxylate,

N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,

N-(4-phenyl-1,3-thiazol-2-yl)-4-propylbenzenesulfonamide,

N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,

N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,

3-Chloro-2-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,

3-Chloro-2-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,

3-Chloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,

2,4,6-Trichloro-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,

2,4,6-Trichloro-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,

- 2,4,6-Trichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-(4-phenyl-1,3-thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
5 N-[4-(4-Fluoro-3-methylphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
2,4-Dichloro-6-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4-Dichloro-6-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4-Dichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-6-
0 methylbenzenesulfonamide,
2,4-Dichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
N-[4-(2-{{(4-propylphenyl)sulfonyl}amino}-1,3-thiazol-4-yl)phenyl]acetamide,
4-Propyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
15 N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-
propylbenzenesulfonamide,
N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-{{(3-chloro-2-methylphenyl)sulfonyl}amino}-1,3-thiazol-4-
20 yl)phenyl]acetamide,
3-Chloro-2-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-2-
methylbenzenesulfonamide,
3-Chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-
25 methylbenzenesulfonamide,
3-Chloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
3-Chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
N-[4-(2-{{(2,4,6-trichlorophenyl)sulfonyl}amino}-1,3-thiazol-4-yl)phenyl]acetamide,
2,4,6-Trichloro-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
30 2,4,6-Trichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,

- 2,4,6-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 5 N-(4-{2-[(1,1'-biphenyl)-4-ylsulfonyl]amino}-1,3-thiazol-4-yl)phenyl)acetamide,
N-[4-(3-pyridinyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- 0 N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(2-{[(2,4-dichloro-6-methylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl)acetamide,
- 2,4-Dichloro-6-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 5 2,4-Dichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 0 2,4-Dichloro-N-[4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- N-[4-(1-benzothien-3-yl)-1,3-thiazol-2-yl]-2,4-dichloro-6-methylbenzenesulfonamide,
- N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- 5 3-Chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
-) Ethyl 2-[(1,1'-biphenyl)-4-ylsulfonyl]amino]-4-phenyl-1,3-thiazole-5-carboxylate,
- 3-Chloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,

- N-[4-(2-[[4-Bromo-2,5-difluorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
- 2,3,4-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,5-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 5 2,3,4-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 4-Bromo-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2,5-difluorobenzenesulfonamide,
- 4,5-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
- 4,5-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 10 N-[4-(2-[[2,4,5-Trichlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
- 2,3,4-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 4-Bromo-5-chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 15 3-Bromo-5-chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
- 3-Bromo-5-chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- N-[4-(2-[[2,6-Dichlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
- 2,6-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 20 2,4,6-Trichloro-N-(7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 2,3,4-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 25 2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- N-[4-(2-Chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- 4-Bromo-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
- 4-Bromo-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
- 30 difluorobenzenesulfonamide,

- 4,5-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 4-Bromo-5-chloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 5 4-Bromo-5-chloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 10 4-Bromo-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 5 N-[4-(2-{{(4-Bromo-5-chloro-2-thienyl)sulfonyl}amino})-1,3-thiazol-4-yl]phenyl]acetamide,
- 3-Chloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 0 N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- 2,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-6-methylbenzenesulfonamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-4-propylbenzenesulfonamide,
- 5 3-Chloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-2-methylbenzenesulfonamide,
- N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-4-propylbenzenesulfonamide,
- 2,4-Dichloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-6-methylbenzenesulfonamide,
-)

N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,

N-(2-{{[(3-Chloro-2-methylphenyl)sulfonyl]amino}naphtho[1,2-d][1,3]thiazol-6-yl)acetamide,

5 N-(2-{{[(4-Propylphenyl)sulfonyl]amino}-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl)acetamide,

N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide,

N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide.


10 4. A compound according to claim 1-3, for medical use.

5. A process for the preparation of a compound according to claim 1-3 comprising at least one of the following steps:

- 15 a) sulfonamide coupling by reacting a 2-aminothiazole with a sulfonylchloride in the presence of a base,
- b) sulfonamide coupling by reacting a 2-aminothiazole derivative with a sulfonylchloride in the presence of a base,
- c) formation of a thiazole ring by reacting an optionally substituted thiourea with an α -haloketone,
- 20 d) formation of a thiazole ring by reacting a thiourea with a ketone.

6. A method for the treatment or prevention of diabetes, syndrome X, obesity, glaucoma, hyperlipidemia, hyperglycemia, hyperinsulinemia, osteoporosis, tuberculosis, depression, virus diseases and inflammatory disorders, said method

25 comprising administering to a mammal, including man, in need of such treatment an effective amount of a compound of the formula (I)



wherein

T is an aryl ring or heteroaryl ring, optionally independently substituted by $[R]_n$,
 5 wherein n is an integer 0-5, and R is hydrogen, halogen, C_{1-6} -alkyl, and aryl;

A is selected from an aryl ring or heteroaryl ring, which can further be optionally substituted in one or more positions independently of each other by hydrogen, C_{1-6} -alkyl, halogenated C_{1-6} -alkyl, halogen, C_{1-6} -alkoxy, nitro, C_{1-6} -alkoxycarbonyl, C_{1-6} -alkylsulfonyl, acetylamino or aryloxy, wherein the aryloxy can further be optionally substituted in one or more positions independently of each other by hydrogen and halogen;
 10

B is selected from hydrogen and C_{1-6} -alkoxycarbonyl or is linked to A to give a 6-
 15 membered aromatic or non-aromatic ring;

as well as pharmaceutically acceptable salts, hydrates and solvates thereof.

7. A method according to claim 6,
 20

wherein

T is selected from
 thienyl substituted with one or more of bromo, chloro;
 25 phenyl optionally substituted with one or more of chloro, methyl, propyl, phenyl, bromo, fluoro;

A is selected from 1-benzothien-3-yl, 3-(2,5-dimethylfuryl), pyridinyl;
thienyl optionally substituted with one or more of chloro, methylsulfonyl;
phenyl optionally substituted with one or more of ethoxycarbonyl, nitro, fluoro,
methyl, methoxy, acetylamino, chloro, 4-chlorophenoxy, trifluoromethyl;

5

B is selected from hydrogen, carbethoxy or is linked to A to give a 6-membered
aromatic or non-aromatic ring.

8. A method according to claim 6-7, wherein the compound is selected from:

- 10 Ethyl 2-(2-{[(4-chlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl)benzoate,
2,5-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
4-Chloro-N-[4-(4,5-dichloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
Ethyl 2-{[(4-chlorophenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-carboxylate,
Ethyl 2-{[(3-chloro-2-methylphenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-
15 carboxylate,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-(4-phenyl-1,3-thiazol-2-yl)-4-propylbenzenesulfonamide,
N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
20 3-Chloro-2-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-2-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
3-Chloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-2-
methylbenzenesulfonamide,
2,4,6-Trichloro-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
5 2,4,6-Trichloro-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4,6-Trichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-(4-phenyl-1,3-thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
0 N-[4-(4-Fluoro-3-methylphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,

- 2,4-Dichloro-6-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4-Dichloro-6-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4-Dichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 5 2,4-Dichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
N-[4-(2-([(4-propylphenyl)sulfonyl]amino)-1,3-thiazol-4-yl)phenyl]acetamide,
4-Propyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-
- 10 propylbenzenesulfonamide,
N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-([(3-chloro-2-methylphenyl)sulfonyl]amino)-1,3-thiazol-4-yl)phenyl]acetamide,
- 15 3-Chloro-2-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
3-Chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-methylbenzenesulfonamide,
- 20 3-Chloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
3-Chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
N-[4-(2-([(2,4,6-trichlorophenyl)sulfonyl]amino)-1,3-thiazol-4-yl)phenyl]acetamide,
2,4,6-Trichloro-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 25 2,4,6-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
2,4,6-Trichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-(4-{2-([(1,1'-biphenyl]-4-ylsulfonyl)amino]-1,3-thiazol-4-yl}phenyl)acetamide,
- 0 N-[4-(3-pyridinyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,


- N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- 5 N-[4-(2-((2,4-dichloro-6-methylphenyl)sulfonyl)amino)-1,3-thiazol-4-yl)phenyl]acetamide,
- 2,4-Dichloro-6-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 10 2,4-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-[4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 5 N-[4-(1-benzothien-3-yl)-1,3-thiazol-2-yl]-2,4-dichloro-6-methylbenzenesulfonamide,
- N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- 3-Chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 0 2,4-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- Ethyl 2-(((1,1'-biphenyl)-4-ylsulfonyl)amino)-4-phenyl-1,3-thiazole-5-carboxylate,
- 3-Chloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 5 N-[4-(2-((4-Bromo-2,5-difluorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl)phenyl]acetamide,
- 2,3,4-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,5-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,3,4-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 1) 4-Bromo-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2,5-difluorobenzenesulfonamide,

- 4,5-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
- 4,5-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- N-[4-(2-[(2,4,5-Trichlorophenyl)sulfonyl]amino)-1,3-thiazol-4-yl]phenyl]acetamide,
- 5 2,3,4-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 4-Bromo-5-chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 3-Bromo-5-chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
- 10 3-Bromo-5-chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- N-[4-(2-[(2,6-Dichlorophenyl)sulfonyl]amino)-1,3-thiazol-4-yl]phenyl]acetamide,
- 2,6-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,6-Trichloro-N-(7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 15 2,3,4-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 20 N-[4-(2-Chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- 4-Bromo-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
- 4-Bromo-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
- 5 4,5-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 4-Bromo-5-chloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 4-Bromo-5-chloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 0

- 2,4-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 5 4-Bromo-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- N-[4-(2-{[(4-Bromo-5-chloro-2-thienyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide,
- 10 N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 3,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 3-Chloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-2-methylbenzenesulfonamide,
- 15 2,4,6-Trichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- 2,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-6-methylbenzenesulfonamide,
- 20 N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-4-propylbenzenesulfonamide,
- 3-Chloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-2-methylbenzenesulfonamide,
- N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-4-propylbenzenesulfonamide,
- 25 2,4-Dichloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-6-methylbenzenesulfonamide,
- N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- N-{2-[(Phenylsulfonyl)amino]-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide,
- 30 N-(2-{[(3-Chloro-2-methylphenyl)sulfonyl]amino}naphtho[1,2-d][1,3]thiazol-6-yl)acetamide,

- N-(2-{[(4-Propylphenyl)sulfonyl]amino}-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl)acetamide,
 N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide,
 N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
 5 N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide.

9. The use of a compound of the formula (I)



0 wherein

T is an aryl ring or heteroaryl ring, optionally independently substituted by [R]_n, wherein n is an integer 0-5, and R is hydrogen, halogen, C₁₋₆-alkyl and aryl;

- 5 A is selected from an aryl ring or heteroaryl ring, which can further be optionally substituted in one or more positions independently of each other by hydrogen, C₁₋₆-alkyl, halogenated C₁₋₆-alkyl, halogen, C₁₋₆-alkoxy, nitro, C₁₋₆-alkoxycarbonyl, C₁₋₆-alkylsulfonyl, acetylamino or aryloxy, wherein the aryloxy can further be optionally substituted in one or more positions independently of each other by hydrogen and
) halogen;

B is selected from hydrogen and C₁₋₆-alkoxycarbonyl or is linked to A to give a 6-membered aromatic or non-aromatic ring;

as well as pharmaceutically acceptable salts, hydrates and solvates thereof,

in the manufacture of a medicament for the treatment or prevention of diabetes, syndrome X, obesity, glaucoma, hyperlipidemia, hyperglycemia, hyperinsulinemia, osteoporosis, tuberculosis, depression, virus diseases and inflammatory disorders.

5 10. The use according to claim 9,

wherein

T is selected from

0 thienyl substituted with one or more of bromo, chloro;
phenyl optionally substituted with one or more of chloro, methyl, propyl, phenyl,
bromo, fluoro;

A is selected from 1-benzothien-3-yl, 3-(2,5-dimethylfuryl), pyridinyl;

5 thienyl optionally substituted with one or more of chloro, methylsulfonyl;
phenyl optionally substituted with one or more of ethoxycarbonyl, nitro, fluoro,
methyl, methoxy, acetylamino, chloro, 4-chlorophenoxy, trifluoromethyl;

B is selected from hydrogen, carbethoxy or is linked to A to give a 6-membered
0 aromatic or non-aromatic ring.

11. The use according to claim 9-10, wherein the compound is selected from:

Ethyl 2-(2-{{[(4-chlorophenyl)sulfonyl]amino}-1,3-thiazol-4-yl})benzoate,
2,5-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
5 4-Chloro-N-[4-(4,5-dichloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
Ethyl 2-{{[(4-chlorophenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-carboxylate,
Ethyl 2-{{[(3-chloro-2-methylphenyl)sulfonyl]amino}-4-phenyl-1,3-thiazole-5-
carboxylate,
N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
10 N-(4-phenyl-1,3-thiazol-2-yl)-4-propylbenzenesulfonamide,
N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,

- N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
3-Chloro-2-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-2-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
3-Chloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-2-
5 methylbenzenesulfonamide,
2,4,6-Trichloro-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4,6-Trichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
10 N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-(4-phenyl-1,3-thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Fluoro-3-methylphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
2,4-Dichloro-6-methyl-N-[4-(3-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
5 2,4-Dichloro-6-methyl-N-(4-phenyl-1,3-thiazol-2-yl)benzenesulfonamide,
2,4-Dichloro-N-[4-(4-fluoro-3-methylphenyl)-1,3-thiazol-2-yl]-6-
methylbenzenesulfonamide,
2,4-Dichloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
N-[4-(2-{{[(4-propylphenyl)sulfonyl]amino}-1,3-thiazol-4-yl)phenyl]acetamide},
0 4-Propyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-
propylbenzenesulfonamide,
N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
5 N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
N-[4-(2-{{[(3-chloro-2-methylphenyl)sulfonyl]amino}-1,3-thiazol-4-
yl)phenyl]acetamide},
3-Chloro-2-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
3-Chloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-2-
0 methylbenzenesulfonamide,

- 3-Chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-methylbenzenesulfonamide,
- 3-Chloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 3-Chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 5 N-[4-(2-((2,4,6-trichlorophenyl)sulfonyl)amino)-1,3-thiazol-4-yl]phenyl]acetamide,
- 2,4,6-Trichloro-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,6-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 10 2,4,6-Trichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4,6-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- N-(4-{2-(((1,1'-biphenyl)-4-ylsulfonyl)amino)-1,3-thiazol-4-yl}phenyl)acetamide,
- N-[4-(3-pyridinyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- 15 N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,
- N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
- N-[4-(2-((2,4-dichloro-6-methylphenyl)sulfonyl)amino)-1,3-thiazol-4-yl]phenyl]acetamide,
- 20 2,4-Dichloro-6-methyl-N-[4-(3-pyridinyl)-1,3-thiazol-2-yl]benzenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chloro-5-nitrophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-6-methylbenzenesulfonamide,
- 5 2,4-Dichloro-N-[4-(5-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 2,4-Dichloro-N-[4-(2,5-dimethyl-3-furyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 0 N-[4-(1-benzothien-3-yl)-1,3-thiazol-2-yl]-2,4-dichloro-6-methylbenzenesulfonamide,
- N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,

- 3-Chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
2,4,6-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
- 5 2,4-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-6-methylbenzenesulfonamide,
Ethyl 2-[[[1,1'-biphenyl]-4-ylsulfonyl]amino]-4-phenyl-1,3-thiazole-5-carboxylate,
3-Chloro-N-[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
N-[4-(2-[[[4-Bromo-2,5-difluorophenyl)sulfonyl]amino]-1,3-thiazol-4-yl)phenyl]acetamide,
- 0 2,3,4-Trichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,5-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,3,4-Trichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
4-Bromo-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2,5-difluorobenzenesulfonamide,
4,5-Dichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
- 5 4,5-Dichloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
N-[4-(2-[[[2,4,5-Trichlorophenyl)sulfonyl]amino]-1,3-thiazol-4-yl)phenyl]acetamide,
2,3,4-Trichloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
- 0 4-Bromo-5-chloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
3-Bromo-5-chloro-N-(7-methoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-2-thiophenesulfonamide,
3-Bromo-5-chloro-N-[4-(2-chlorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 5 N-[4-(2-[[[2,6-Dichlorophenyl)sulfonyl]amino]-1,3-thiazol-4-yl)phenyl]acetamide,
2,6-Dichloro-N-[4-(3-chloro-2-thienyl)-1,3-thiazol-2-yl]benzenesulfonamide,
2,4,6-Trichloro-N-(7,8-dimethoxy-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
2,3,4-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 1

- 2,3,4-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- N-[4-(2-Chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-4-propylbenzenesulfonamide,
- 4-Bromo-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2,5-
- 5 difluorobenzenesulfonamide,
- 4-Bromo-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2,5-difluorobenzenesulfonamide,
- 4,5-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-thiophenesulfonamide,
- 0 4-Bromo-5-chloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 4-Bromo-5-chloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}-2-thiophenesulfonamide,
- 2,4-Dichloro-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-6-
- 5 methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-{4-[2,6-dichloro-4-(trifluoromethyl)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- 4-Bromo-N-[4-(2-chloro-6-fluorophenyl)-1,3-thiazol-2-yl]-2-methylbenzenesulfonamide,
- 0 2,4,6-Trichloro-N-{4-[2-chloro-4-(4-chlorophenoxy)phenyl]-1,3-thiazol-2-yl}benzenesulfonamide,
- N-[4-(2-{{(4-Bromo-5-chloro-2-thienyl)sulfonyl}amino}-1,3-thiazol-4-yl)phenyl]acetamide,
- N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 5 3,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
- 3-Chloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-2-methylbenzenesulfonamide,
- 2,4,6-Trichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)benzenesulfonamide,
-) N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide,

- 2,4-Dichloro-N-(4,5-dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-6-methylbenzenesulfonamide,
N-(4,5-Dihydrothieno[3,2-e][1,3]benzothiazol-2-yl)-4-propylbenzenesulfonamide,
3-Chloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-2-methylbenzenesulfonamide,
5 N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-4-propylbenzenesulfonamide,
2,4-Dichloro-N-[6-chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl]-6-methylbenzenesulfonamide,
0 N-[6-Chloro-8-(methylsulfonyl)-4,5-dihydrothieno[3,4-e][1,3]benzothiazol-2-yl][1,1'-biphenyl]-4-sulfonamide,
N-{2-[(Phenylsulfonyl)amino]-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl}acetamide,
N-(2-[(3-Chloro-2-methylphenyl)sulfonyl]amino)naphtho[1,2-d][1,3]thiazol-6-yl)acetamide,
5 N-(2-[(4-Propylphenyl)sulfonyl]amino)-4,5-dihydronaphtho[1,2-d][1,3]thiazol-6-yl)acetamide,
N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)-4-propylbenzenesulfonamide,
N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)benzenesulfonamide,
N-(8-Nitro-4,5-dihydronaphtho[1,2-d][1,3]thiazol-2-yl)[1,1'-biphenyl]-4-sulfonamide.
0
12. A pharmaceutical composition comprising at least one compound of the formula (I) as defined in claim 1-3, and a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/01158

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 277/52, C07D 277/60, A61K 31/426, A61P 3/00, A61P 5/48, A61P 27/06,
A61P 25/24, A61P 25/28, A61P 29/00, A61P 31/12, A61P 31/06
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CHEM.ABS DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 94123 E (DR. KARL THOMAE G.M.B.H.), 27 May 1969 (27.05.69) --	1-12
X	NL 6610324 A (DR. KARL THOMAE GESELLSCHAFT MIT BESCHRÄNKTER HAFTUNG), 24 January 1967 (24.01.67) --	1-12
X	Chemical Abstracts, Volume 59 (6), (Columbus, Ohio, USA), J. D. McColl et al, "Effect of sulfonylurea derivatives in experimental ulcer formation in the rat", THE ABSTRACT No 3231, Arch. Intern. Pharmacodyn. 1963, 141, 181-189 --	1-12

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier application or patent but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

Date of the actual completion of the international search

4 Sept 2001

Date of mailing of the international search report

11-09-2001

Name and mailing address of the ISA/
 Swedish Patent Office
 Box 5055, S-102 42 STOCKHOLM
 Facsimile No. +46 8 666 02 86

Authorized officer
 Gerd Strandell/EÖ
 Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/01158

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STN International, File CAPLUS, CAPLUS accession no. 1991:680016, Document no. 115:280016, Hisamitsu Pharmaceutical Co.: "Preparation of diphenylthiazoles as pharmaceuticals"; JP,A2,03173876, 19910729 --	1-12
X	WO 9928306 A1 (PHARMACIA & UPJOHN S.P.A.), 10 June 1999 (10.06.99) --	1-12
X	EP 0819681 A2 (F. HOFFMANN-LA ROCHE AG), 21 January 1998 (21.01.98) --	1-12
X	EP 0790057 A1 (HISAMITSU PHARMACEUTICAL CO. INC.), 20 August 1997 (20.08.97) --	1-12
X	STN International, File CAPLUS, CAPLUS accession no. 1995:867676, Document no. 123:256699, Hisamitsu Pharmaceutical Co: "Preparation of 2-aminothiazole derivatives as esterase inhibitors"; JP,A2,07149745, 19950613 --	1-12
X	STN International, File CAPLUS, CAPLUS accession no. 1995:818696, Document no. 123:228174, Hisamitsu Pharmaceutical Co: "Preparation of 2-(substituted amino)thiazole derivatives as esterase inhibitors"; & JP,A2,07149746, 19950613 --	1-12
X	Chemical Abstracts, Volume 59, (Columbus, Ohio, USA), Gaile, E. Gudriniece et al: "Heterocyclic compounds based on diketones. II. 2'-Amino-5,5-dimethyl-1-cyclohexanone(2,3:4',5')thiazole. I.", THE ABSTRACT No 6380, Latvijas PSR Zinatnu Akad. Vestis., Kim. Ser. 1962, No. 4, 523-8 --	1-3,5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/01158

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9707789 A1 (THE UNIVERSITY OF EDINBURGH), 6 March 1997 (06.03.97) -- -----	1-12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE01/01158

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 6-8
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE01/01158

Claims 6-8 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

INTERNATIONAL SEARCH REPORT
Information on patent family members

02/08/01

International application No.

PCT/SE 01/01158

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
FR	94123	E	27/05/69	ES	349314 A	01/04/69
				NL	6800598 A	22/07/68
				ES	349316 A	01/04/69
				NL	6800589 A	17/07/68
				ES	349317 A	01/04/69
				ES	349319 A	01/04/69
				ES	349320 A	01/04/69
				ES	349321 A	01/04/69
				ES	349322 A	01/04/69
				ES	349318 A	01/04/69
NL	6610324	A	24/01/67	BE	684415 A	20/01/67
				BR	6681517 D	00/00/00
				DE	1620508 A	18/09/69
				ES	329299 A	16/02/68
				FR	6013 M	06/05/68
				FR	1498465 A	08/01/68
				GB	1140387 A	15/01/69
WO	9928306	A1	10/06/99	AU	1753599 A	16/06/99
				EP	1036069 A	20/09/00
				GB	9725141 D	00/00/00
EP	0819681	A2	21/01/98	AU	725496 B	12/10/00
				AU	2875297 A	29/01/98
				BR	9704023 A	17/11/98
				CA	2210613 A	19/01/98
				CN	1176101 A	18/03/98
				CZ	9702273 A	18/02/98
				HR	970393 A	30/04/98
				HU	9701193 A	01/02/99
				IL	121298 D	00/00/00
				JP	2991679 B	20/12/99
				JP	10067761 A	10/03/98
				KR	247672 B	01/04/00
				NO	973338 A	20/01/98
				NZ	328331 A	28/05/99
				PL	321203 A	02/02/98
				TR	9700653 A	00/00/00
				US	5877193 A	02/03/99
				US	5958910 A	28/09/99
EP	0790057	A1	20/08/97	AU	689972 B	09/04/98
				AU	3880995 A	19/06/96
				JP	3023178 B	21/03/00
				US	5856347 A	05/01/99
				CA	2206315 A	06/06/96
				TW	414708 B	00/00/00
				WO	9616650 A	06/06/96
WO	9707789	A1	06/03/97	AU	6833796 A	19/03/97
				EP	0847275 A	17/06/98
				GB	2317826 A,B	08/04/98
				GB	9517622 D	00/00/00
				GB	9801921 D	00/00/00

Form PCT/ISA.210 (patent family annex) (July 1998)